arXiv:1501.06376v1 [math.PR] 26 Jan 2015

Weak Law of Large Numbers and Central Limit Theorem for probabilistic system with given Entropy

Vassili Kolokoltsov (v.kolokoltsov@warwick.ac.uk),
Tomasz M. Lapinski (84tomek@gmail.com)

University of Warwick,
Department of Statistics,
CV4 7AL Coventry, UK,

January 27, 2015

Abstract

In this paper we introduce Law of Large Numbers and Central Limit Theorem for certain probabilistic system with estimated rate of convergence. The underlying probability space on which results are constructed is determined by the Entropy, which is a number of configurations of system which can occur for given state. The sample space is a lattice of rational numbers and represents the system states. Probability mass function of particular state is obtained by normalizing the Entropy of considered state by the sum of Entropies of all states. The Entropy has a unique maximum for considered lattice and several other technical properties. As the system grows, i.e. its certain parameter increases, the Entropies increases and the lattice becomes denser. The first result of the paper gives the mean state as systems size tends to infinity. The second result, Central Limit Theorem, depends on the type of maximum of entropy. When maximum is in the interior of the lattice the fluctuation converges to Gaussian distribution. In case the maximum is on the boundary of the domain the distribution is a discrete in the direction orthogonal to the boundary of lattice on which maximum is situated and Gaussian in other directions. Additionally, for each result we provide the rate of convergence of the system to the resulting distribution. The proof consists of several approximations developed specially for this paper. The main are the Laplace type approximation but with sum instead of integrals and several other but rather standard approximations.

Key words— law of large numbers, central limit theorem, laplace approximation, entropy

1 Introduction

Limit theorems is a very rich and important branch of the probability theory. The Law of Large numbers and Central Limit theorems are among the fundaments of probabilistic theory and are essential in many probabilistic applications. In this paper we present a new theoretical results for certain probabilistic system.

The states of the system are represented as the nods on some finite lattice of rational numbers. For each nod, i.e. state, we are given an Entropy of the system, i.e. the number of configurations which system can take. The number of states and the Entropy depends on some parameter. As it increases the Entropy of the states increases and the lattice of states becomes more dense, i.e. the distances between nods become smaller. We can vaguely say that our probabilistic system increases as this parameter increase. The Entropy has a unique maximum on considered domain and for all value of parameter. Additionally Entropy has several others properties, rather technical in nature used in the proof.

With this probabilistic system there is associated a formal probability space. The state space are the rational numbers which corresponds to the nodes of the lattice and there is associated with it a discrete sigma algebra. The probability measure, i.e. the probability of some set of states is just the sum exponents of the entropy of those states normalized by the sum of exponents of Entropies of all states.
With the occurrence of certain state on the lattice, i.e. rational number, there is associated a random variable. The first result is a Weak Law of Large number for this random variables. It gives us a single state to which system converges as the systems grows to infinity. This single state is one for which the Entropy has maximal value, and it can be called a mean.
The second result is associated with the random variables which correspond to deviation from the mean obtained from previous result, multiplied by our increasing parameter to some power. It gives us the distribution of deviation of the system from the 'average' state. For two types of maximum the Entropy we obtain different distributions. For the case when maximum is in the interior of the the lattice is we obtain a Gaussian distribution. The second case is when the maximum is on the boundary of the lattice then we have a discrete distribution in direction orthogonal to boundary on which maximum is situated and Gaussian in other directions.

The method of proving of both theorems is similar. We use the fact the weak convergence of random variable is implied by the convergence of the moment generating functions of this random variable. The moment generating functions are composed of sums of Entropies over all states in the denominator and with some additional factor also in the numerator. We developed a result for approximating those sums. Essentially, it is similar to Multivariate Laplace approximation introduced in Kolokoltsov [2000] but is more complex, due to the summation instead of integration. Once the the sums are approximated, some minor additional approximations are needed but are rather standard. Additionally, within the proof there is a special care taken to provide a rate of convergence of the system to given result.

The paper consist of six sections and appendix. The second section of the paper introduces a mathematical setting on which the results are build on. We introduce the lattice, Entropy, its properties and prove its necessary properties. The probability space is also formally introduced. In the third section we introduce and prove our two limit theorems. The fourth section provides Laplace type results. There two results, one for each type of maximum of Entropy. In the next section we contained result used in the Laplace type approximation, its rather standard. It is a approximation of Gaussian integral with the sum. Finally, in the last section we put an additional estimate which are rather standard but developed specially for this paper. In the appendix we put relatively standard results used in the proofs.

2 Mathematical setting

We consider a function $S : \Omega_N \times \mathbb{N} \to \mathbb{R}$ which has a unique maximum at the point $x^*(N)$, for every $N > N_0$. In the limit, as $N \to \infty$, the unique maximums tend to the point $x^*(N) \to x^*$.

Let Ω be some convex subset of $\mathbb{R}^m$. Then let us consider the lattice of the points $x \in \mathbb{R}^m$ separated by a distance $\frac{1}{N}$, i.e. their components x_i, $i, \ldots, m$ takes values from the set $\{0, \frac{1}{N}, \frac{2}{N}, \frac{3}{N}, \ldots, 1, \frac{N+1}{N}, \frac{N+2}{N}, \ldots\}$. The domain of the function S, the set Ω_N for considered N is a subset of that lattice such that it consists of the points contained in the subset Ω.

Then we define a probability space. The domain Ω_N is the sample space and the events given by x's are its points on the considered lattice. Further we define a m-dimensional discrete random vector $X_N : \Omega_N \to \mathbb{R}^m$ with pmf

$$P(X_N = x) = \frac{e^{S(x,N)}}{\sum_{\Omega_N} S(x,N)}. \tag{1}$$

Moreover, for the function S we consider two types of maximum

a) Maximum inside the domain separated from the boundary by at least some constant distance in the limit as $N \to \infty$

b) Maximum at the boundary for some $N > N_0$ and the point of maximum is not a critical. This boundary is a subset of a hyperplane in $\mathbb{R}^m$. The number v_1^* is the first component of $v^*(N)$ defined by $Tv^*(N) = x^*(N)$. Transformation T is rotation of coordinate system such that v_1^* represents a distance form the origin to the boundary in appropriately rotated coordinate system is rational and .

The function $S(x,N)$, for large enough N is such that

$$DS(x,N) = h(N)[Ds(x) + \sigma(x)\epsilon(N)], \tag{2}$$

where D is a differential operator and $\epsilon(N) \to 0$ as $N \to \infty$. Functions s and σ are C^2 functions, are three times differentiable and matrix D^2S is negative definite for all $x \in \Omega$ and some large enough N. Further the function $h(N) = N$ or is such that $\frac{h(N)}{N} \to 0$. The resulting properties of the function S are given by the lemma

Lemma 1. *For the function S we have constants $s^{(2)}, s'^{(2)}$ and $s^{(3)}$ such that*

$$s^{(2)} = \sup_{x \in \Omega_N, N > N_0} \frac{1}{h(N)} \|D^2 S(x,N)\| < \infty, \tag{3}$$

Since Hessian $D^2 s(x)$ is invertible we have

$$s'^{(2)} = \inf_{x \in \Omega_N, N > N_0} \frac{1}{h(N)} \|D^2 S(x,N)^{-1}\|^{-1} > 0, \tag{4}$$

and

$$s^{(3)} = \sup_{x \in \Omega_N, N > N_0} \frac{1}{h(N)} \|D^{\otimes 3} S(x,N)\| < \infty. \tag{5}$$

The point of maximum of $S(x,N)$ is asymptotically equal to

$$x^*(N) = x^* + O(1)\epsilon(N), N \to \infty \tag{6}$$

Proof. We start by proving the first result, (3). We calculate the second derivative of the representation of derivative vector DS given by (2)

$$D^2S(x,N) = h(N)[D^2s(x) + D\sigma(x)\epsilon(N)], \tag{7}$$

where $\sigma(x)$ is a vector where each component cooreponds to a component in vector x. Now we take the norm, apply triangle inequality and divide by $h(N)$ and get

$$\frac{1}{h(N)}\|D^2S(x,N)\| \leq \|D^2s(x)\| + \epsilon(N)\|D\sigma(x)\|.$$

Since the first expression is a constant for any $x \in \Omega_N$ and the second one tends to 0 as $N \to \infty$, hence LHS is bounded for $N > N_0$ and $x \in \Omega_N$ and therefore (3) is valid.
Now, let us prove (4). We calculate the inverse of the second derivative D^2S (7)

$$D^2S(x,N)^{-1} = \frac{D^2s(x)^{-1}}{h(N)}\Bigg[I + D^2s(x)^{-1}D\sigma(x)\epsilon(N)\Bigg]^{-1},$$

which is possible since D^S is negative definite, hence invertible.
Now we apply the asymptotic version of second result of the Theorem 5 from the Appendix

$$(I + \epsilon A)^{-1} = I + O(\epsilon),\ \epsilon \to 0, \tag{8}$$

and obtain

$$D^2S(x,N)^{-1} = \frac{D^2s(x)^{-1}}{h(N)}[I + O(\epsilon(N))],$$

Then we take a norm, put expression in denominator and divide by $h(N)$

$$\frac{1}{h(N)}\|D^2S(x,N)^{-1}\|^{-1} = \|D^2s(x)^{-1}[I + O(\epsilon(N))]\|^{-1},$$

and the apply Schwartz inequality and again result (8)

$$\frac{1}{h(N)}\|D^2S(x,N)^{-1}\|^{-1} \geq \|D^2s(x)^{-1}\|^{-1}\|I + O(\epsilon(N))\|,$$

Since first expression on the RHS is constant for all $x \in \Omega_N$ and second is bounded for all N, hence LHS is bounded hence we get the final result. For the last result (5) we calculate the third derivatives tensor $D^{\otimes 3}S$, take the norm, apply triangle inequality and divide by $h(N)$ ad obtain

$$\frac{1}{h(N)}\|D^{\otimes 3}S(x,N)\| \leq \|D^{\otimes 3}s(x)\| + \epsilon(N)\|D^2\sigma(x)\|.$$

Again, since all the expressions on the RHS are bounded the LHS is also bounded for $N > N_0$ and $x \in \Omega_N$, hence we proved the result.
We begin proving the last result for maximum of type a) with first order Taylor expansion of the vector $DS(x^*(N),N)$ at the point x^*

$$DS(x^*(N),N) = DS(x^*,N) + D^2S(x_{\theta_1}(N),N)(x^*(N) - x^*) \tag{9}$$

where $x_{\theta_1}(N)$ is some vector between $x^*(N)$ and x^*. Since $x^*(N)$ is a maximum of $S(x,N)$ and the maximum is of the type a) we have $DS(x^*(N),N) = o$. Further, since x^* is a maximum in the limit hence using representation (2) $\lim_{N\to\infty}[Ds(x^*)+\sigma(x^*)\epsilon(N)]=o$ we get $DS(x^*,N)=h(N)\sigma(x^*)\epsilon(N)$. Then (9) is equivalent

$$o=h(N)\sigma(x^*)\epsilon(N)+D^2S(x_{\theta_1}(N),N)(x^*(N)-x^*) \tag{10}$$

and after taking absolute values and dividing by the norm of D^2S

$$|h(N)\sigma(x^*)\epsilon(N)|=\|D^2S(x_{\theta_1}(N),N)\||x^*(N)-x^*|$$

and then we bound the norm using formula form appendix (80) and property of S (4)

$$|h(N)\sigma(x^*)\epsilon(N)|=\|D^2S(x_{\theta_1}(N),N)\||x^*(N)-x^*|\geq h(N)s'^{(2)}|x^*(N)-x^*|$$

hence finally we have a bound

$$\frac{1}{s'^{(2)}}|\sigma(x^*)\epsilon(N)|\geq|x^*(N)-x^*|$$

which is valid for large enough N since we used representation (2) and is equivalent to the result when put in the big 'O' notation. This was result for the maximum of type a). For the maximum of typa b) we have to perform the change of coordinates $x=Tv$ first and than as previously use Taylor theorem

$$DS(Tv^*(N),N)=DS(Tv^*,N)+D^2S(Tv_{\theta_1}(N),N)(Tv^*(N)-Tv^*) \tag{11}$$

Since the maximum is on the boundary, which is orthogonal to the direction v_1, therefore $v_1^*(N)=v_1^*$ is the same for all N in the vector $v^*(N)$. In other directions the maximum is varying and is in the point for which the first derivatives are equal to 0. Therefore the result (10) is the same for maximum of type b) with the difference that the vector $h(N)\sigma(x^*)\epsilon(N)$ has first component equal to 0, i.e.

$$\begin{pmatrix}0\\ h(N)\hat{\sigma}(x^*)\epsilon(N)\end{pmatrix}=D^2S(x_{\theta_1}(N),N)(x^*(N)-x^*)$$

where $\hat{\sigma}(x^*)$ is a $m-1$ dimensional vector.
We have this 0 because the first derivatives in (11) cancel out for the first component as $v_1^*(N)=v_1^*$. Then we perform the same calculations as for the maximum type a) and the difference in the final result lays only in the constant hence asymptotic result is the same as for maximum of type a). ☐

3 Law of large numbers and corresponding fluctuations theorem

Theorem 1 (Weak Law of large numbers). *Let X_N be the m-dimensional discrete random vector on the sample space Ω_N with pmf specified by (1). As $N\to\infty$ the random vector X_N converges in distribution to the constant vector $x^*=(x_1^*,x_2^*,\dots,x_m^*)$ and we have following estimates, the same for both types of the maximum*

$$M_{X_N}(\xi)=e^{\xi^Tx^*}+O\left(\frac{1}{h(N)^{1/2-3\delta}}\right),\qquad \textit{when } \frac{1}{h(N)^{1/2-3\delta}}\gg\epsilon(N),$$
$$M_{X_N}(\xi)=e^{\xi^Tx^*}+O(\epsilon(N)),\qquad \textit{when } \frac{1}{h(N)^{1/2-3\delta}}\ll\epsilon(N),$$

where $\delta \in (0, \min\{1/2m, 1/6\})$ *is a constant,* $N \to \infty$*,* $M_{X_N}(\xi)$ *is a moment generating function of the random vector* X_N*.*

Proof. We prove the theorem by showing convergence of the moment generating function of the random vector X_N to a constant vector x^* as $N \to \infty$.

The mgf of r.v. X_N is equal

$$M_{X_N}(\xi) = E[e^{\xi^T X}].$$

Evaluating the probability mass function we obtain following expression for $M_{X_N}(\xi)$

$$M_X(\xi) = \frac{\sum_{\Omega_N} e^{\xi^T x} e^{S(x,N)}}{\sum_{\Omega_N} e^{S(x,N)}}. \tag{12}$$

We start with approximating the first part of $M_{X_N}(\xi)$, i.e the denominator for the two cases a) and b) using the Lemma 2 form the Section 2

$$a) \sum_{\Omega_N} e^{S(x,N)} = \frac{N^m}{d} e^{S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}} \left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right),$$

$$b) \sum_{\Omega_N} e^{S(x,N)} = \frac{N^{m-1}}{d} e^{S(x^*(N),N)} \frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}} \sum_{i=0}^{\infty} e^{\frac{1}{N} S'(x^*(N),N) i} \left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right),$$

where in this case the function $g = 1$ in the Lemma and we change the form of the estimates. Analogically we approximate the numerator of the mgf (12). The additional function under the sums simply our function g in the estimates

$$a) \sum_{\Omega_N} e^{\xi^T x + S(x,N)} = \frac{N^m}{d} e^{\xi^T x^*(N) + S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}} \left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right),$$

$$b) \sum_{\Omega_N} e^{\xi^T x + S(x,N)} = \frac{N^{m-1}}{d} e^{\xi^T x^*(N) + S(x^*(N),N)} \frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}} \times$$
$$\times \sum_{i=0}^{\infty} e^{\frac{1}{N} S'(x^*(N),N) i} \left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right),$$

Then we put together the approximations of the numerator and denominator of mgf using Lemma 7 from the Appendix, cancel the identical terms and for both types of maximum we have

$$M_{X_N}(\xi) = e^{\xi^T x^*(N)} \left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right),$$

as $N \to \infty$.
Next we use following Taylor expansion

$$e^{\xi^T x^*(N)} = e^{\xi^T x^*} + \xi e^{\xi^T x_\theta(N)} (x^*(N) - x^*),$$

where $x_\theta(N)$ is a point between $x^*(N)$ and x^*. Then we substitute the estimate for $x^*(N) - x^*$ (6) and obtain

$$e^{\xi^T x^*(N)} = e^{\xi^T x^*} + \xi e^{\xi^T x_\theta(N)} \epsilon(N),$$

and we substitute it into (13) and get an estimate

$$M_{X_N}(\xi) = \left[e^{\xi^T x^*} + \xi e^{\xi^T x_\theta(N)}\epsilon(N)\right]\left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right),$$

hence we obtain two cases depending which term is asymptotically 'stronger' $\epsilon(N)$ or $\frac{1}{h(N)^{1/2-3\delta}}$

$$\begin{aligned} M_{X_N}(\xi) &= e^{\xi^T x^*} + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right), && \text{when } \frac{1}{h(N)^{1/2-3\delta}} \gg \epsilon(N), \\ M_{X_N}(\xi) &= e^{\xi^T x^*} + O(\epsilon(N)), && \text{when } \frac{1}{h(N)^{1/2-3\delta}} \ll \epsilon(N), \end{aligned}$$

where $\delta \in (0, \min\{1/2m, 1/6\})$ is a constant and $N \to \infty$

□

For the discrete random vector X_N on the sample space Ω_N with pmf specified by (1) we have a m-dimensional random vector Y_N separately for the maximum of type a) and b)

$$\begin{aligned} &a)\ Y_N = \sqrt{h(N)}(X_N - x^*), \\ &b)\ Y_N = N(V_1 - v_1^*) + \sqrt{N}(\hat{V} - \hat{v}^*), \end{aligned} \tag{13}$$

where x^* and $x^* = Tv^* = T(v_1^*, \hat{v}^*)$ is the maximum of the functions S on Ω_N. The transformation $x = Tv$ is a rotation of the coordinate system such that the axis v_1 is orthogonal to the hyperplane of Ω_N on which maximum is attained. In the new coordinate system we have a transformed random vector $X_N = TV = T(V_1, \hat{V})$.

Theorem 2 (Central Limit Theorem). *As $N \to \infty$ and if $\epsilon(N) \ll 1/\sqrt{h(N)}$ from the representation (2) the defined above random vector Y_N converges in distribution to a random vector with normal distribution $\mathcal{N}(0, D^2 s(x^*)^{-1})$ for the maximum type a) and for the maximum of type b) to the mixture of normal distribution $\mathcal{N}(0, \hat{D}^2 s(x^*)^{-1})$ along variable $\hat{V}$ and discrete distribution with pmf $\frac{\exp\{is'(x^*)\}}{\sum_{i=1}^{\infty}\exp\{is'(x^*)\}}$, where i is an index of points in the coordinate v_1 on the lattice $T(\Omega_N)$ starting form the maximal point v_1^*. The matrix of derivatives $\hat{D}^2 s(x^*)^{-1}$ is $m-1 \times m-1$ matrix along v_i, $i = 2, \ldots, m$ and $s'(x^*)$ is derivative of s along v_1. The estimate is valid for a subsequence of integers N which elements can be divided by some integer q, where $\frac{p}{q} = v_1^*$ with $v^* = T^{-1}x^*$. Moreover, we assume that $h(N) = N$ in the representation (2) of S for the second type of maximum. Furthermore, we have estimates*

$$\begin{aligned} &a)\ M_{Y_N}(\xi) = e^{\frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi}\left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right), && \textit{when } \epsilon(N) \ll \frac{1}{h(N)^{1-3\delta}}, \\ &\quad M_{Y_N}(\xi) = e^{\frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi}\left(1 + O\left(\sqrt{h(N)}\epsilon(N)\right)\right), && \textit{when } \epsilon(N) \gg \frac{1}{h(N)^{1-3\delta}}, \\ &b)\ M_{Y_N}(\xi) = \frac{\sum_{i=1}^{\infty} e^{is'(x^*)+\xi_1 i}}{\sum_{i=1}^{\infty} e^{is'(x^*)}} e^{\frac{1}{2}\hat{\xi}^T \hat{D}^2 s(x^*)^{-1}\hat{\xi}}\left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right), && \textit{when } \epsilon(N) \ll \frac{1}{h(N)^{1-3\delta}}, \\ &\quad M_{Y_N}(\xi) = \frac{\sum_{i=1}^{\infty} e^{is'(x^*)+\xi_1 i}}{\sum_{i=1}^{\infty} e^{is'(x^*)}} e^{\frac{1}{2}\hat{\xi}^T \hat{D}^2 s(x^*)^{-1}\hat{\xi}}\left(1 + O\left(\sqrt{h(N)}\epsilon(N)\right)\right), && \textit{when } \epsilon(N) \gg \frac{1}{h(N)^{1-3\delta}}, \end{aligned}$$

as $N \to \infty$, where $\delta \in (0, \min\{1/2m, 1/6\})$ is some arbitrary small constant and M_{Y_N} is a moment generating function of random vector Y_N.

Proof. The proof is analogical to the prove of the previous theorem. We consider the moment generating function of the random vector Y_N

$$M_{Y_N}(\xi) = E[e^{\xi^T Y_N}],$$

and evaluating the probability mass function we obtain following expression for $M_{Y_N}(\xi)$

$$M_{Y_N}(\xi) = \frac{\sum_{\Omega_N} e^{\widetilde{S}(x,N)}}{\sum_{\Omega_N} e^{S(x,N)}}. \tag{14}$$

where $\widetilde{S}(x,N) = S(x,N) + \xi^T Y_N$. Since the Y_N are of the form (13) the function $\widetilde{S}$ has the same properties as S. The corresponding representation (2) for $\widetilde{S}$ are given by

$$a)\ D\widetilde{S}(x,N) = h(N)[Ds(x) + \sigma(x)\widetilde{\delta}(N)],\ \widetilde{\delta}(N) = \epsilon(N) + \frac{1}{\sigma(x)}\frac{1}{\sqrt{h(N)}},$$

$$b)\ D\widetilde{S}(x,N) = h(N)[D\widetilde{s}(x) + \widetilde{\delta}(N)],\ \widetilde{\delta}(N) = \epsilon(N) + \frac{1}{\hat{\sigma}(x)\sqrt{N}},\ \ s'(x) = s'(x) + 1$$

and for $\widetilde{S}$ two types of maximum are not altered.
We first prove the result for the maximum of type a). We approximate the numerator and denominator of the mgf (14) using Lemma 2 and combine them using Lemma 7. Then cancel identical terms and obtain

$$M_{Y_N}(\xi) = e^{\widetilde{S}(\widetilde{x}^*(N),N) - S(x^*(N),N)} \frac{\sqrt{\det D^2 S(x^*(N),N)}}{\sqrt{\det D^2 \widetilde{S}(\widetilde{x}^*(N),N)}}\Bigg|\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right).$$

Next we insert the estimate (64) from Preposition 1

$$M_{Y_N}(\xi) = e^{\widetilde{S}(\widetilde{x}^*(N),N) - S(x^*(N),N)}\left(1 + O\left(\frac{1}{\sqrt{N}}\right)\right)\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right),$$

and we simplify the estimate including only the 'strongest' term as $N \to \infty$

$$M_{Y_N}(\xi) = e^{\widetilde{S}(\widetilde{x}^*(N),N) - S(x^*(N),N)}\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right).$$

Finally we substitute the estimate (63) from Preposition 1

$$M_{Y_N}(\xi) = e^{-\frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi + O(1)\frac{1}{\sqrt{h(N)}}}\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right) \qquad \text{when } \epsilon(N) \ll \frac{1}{h(N)},$$

$$M_{Y_N}(\xi) = e^{-\frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi + O(1)\sqrt{h(N)}\epsilon(N)}\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right) \quad \text{when } \epsilon(N) \gg \frac{1}{h(N)}.$$

Since the second terms in the exponent goes to 0 as $N \to \infty$ we can estimate

$$e^{O(1)\frac{1}{\sqrt{h(N)}}} = 1 + O(1)\frac{1}{\sqrt{h(N)}},$$

$$e^{O(1)\sqrt{h(N)}\epsilon(N)} = 1 + O(1)\sqrt{h(N)}\epsilon(N),$$

and we substitute that back into estimate and get

$$M_{Y_N}(\xi) = e^{-\frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi}\left(1 + O(1)\frac{1}{\sqrt{h(N)}}\right)\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right) \qquad \text{when } \epsilon(N) \ll \frac{1}{h(N)},$$
$$M_{Y_N}(\xi) = e^{-\frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi}\left(1 + O(1)\sqrt{h(N)}\epsilon(N)\right)\left(1 + O\left(\frac{1}{N^{1/2-3\delta}}\right)\right) \quad \text{when } \epsilon(N) \gg \frac{1}{h(N)},$$

and then simplifying the estimate and altering appropriately these cases we get the result of the theorem.

For the maximum of type b) we perform similar calculations. Again, we approximate the numerator and denominator of the mgf (14) using this time Lemma 3 and combine them using Lemma 7, cancel identical terms and obtain

$$M_{Y_N}(\xi) = e^{\widetilde{S}(\widetilde{x}^*(N),N)-S(x^*(N),N)}\frac{\sum_{i=1}^{\infty} e^{\frac{1}{N}\widetilde{S}'(\widetilde{x}^*(N),N)}i}{\sum_{i=1}^{\infty} e^{\frac{1}{N}S'(x^*(N),N)i}}\frac{\sqrt{\det \hat{D}^2 S(x^*(N),N)}}{\sqrt{\det \hat{D}^2 \widetilde{S}(\widetilde{x}^*(N),N)}}\left(1+O\left(\frac{1}{N^{1/2-3\delta}}\right)\right).$$

and then we use the results of the Preposition 1 to and obtain estimate

$$M_{Y_N}(\xi) = \frac{\sum_{i=1}^{\infty} e^{\frac{1}{N}\widetilde{S}'(\widetilde{x}^*(N),N)}i}{\sum_{i=1}^{\infty} e^{\frac{1}{N}S'(x^*(N),N)i}} e^{\frac{1}{2}\hat{\xi}^T \hat{D}^2 s(x^*)^{-1}\hat{\xi}}\left(1 + O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right), \quad \text{when } \epsilon(N) \ll \frac{1}{h(N)^{1-3\delta}},$$
$$M_{Y_N}(\xi) = \frac{\sum_{i=1}^{\infty} e^{\frac{1}{N}\widetilde{S}'(\widetilde{x}^*(N),N)}i}{\sum_{i=1}^{\infty} e^{\frac{1}{N}S'(x^*(N),N)i}} e^{\frac{1}{2}\hat{\xi}^T \hat{D}^2 s(x^*)^{-1}\hat{\xi}}\left(1 + O\left(\sqrt{h(N)}\epsilon(N)\right)\right), \quad \text{when } \epsilon(N) \gg \frac{1}{h(N)^{1-3\delta}}, \tag{15}$$

Next we use first order Taylor theorem for the function $\widetilde{S}'(\widetilde{x}^*(N),N)$ at x^*

$$\widetilde{S}'(\widetilde{x}^*(N),N) = \widetilde{S}'(x^*,N) + D\widetilde{S}'(x_\theta(N))(\widetilde{x}^*(N) - x^*)$$

Since the second derivatives $D\widetilde{S}' = DS'$ and because of (3) the norm of vector of second derivatives is also bounded $\|DS'\| \leq h(N)\bar{s}^{(2)}$. Hence we can write above estimate equation as

$$|\widetilde{S}'(\widetilde{x}^*(N),N) - \widetilde{S}'(x^*,N)| \leq h(N)\bar{s}^{(2)}(|\widetilde{x}^*(N) - x^*(N)| + |x^*(N) - x^*|),$$

and then we use estimate (6) and (62) in the simplifiesd form and get

$$|\widetilde{S}'(\widetilde{x}^*(N),N) - \widetilde{S}'(x^*,N)| \leq h(N)\bar{s}^{(2)}(K_1|1/\sqrt{h(N)}| + K_2|\epsilon(N)|).$$

Since we have assumption that $\epsilon(N) \ll \frac{1}{\sqrt{h(N)}}$ we can write above as

$$\widetilde{S}'(\widetilde{x}^*(N),N) = \widetilde{S}'(x^*,N) + O(1)\sqrt{h(N)},$$

and now we use the representation (2) of S

$$\widetilde{S}'(\widetilde{x}^*(N),N) = h(N)(s'(x^*) + \xi_1 + \sigma_1(x^*)\epsilon(N)) + O(1)\sqrt{h(N)},$$

where we used explicit form of $\widetilde{S}$ and then again we use the assumption about $\epsilon(N)$ and since $h(N) = N$ for the maximum of type b) we finally obtain

$$\widetilde{S}'(\widetilde{x}^*(N),N) = N(s'(x^*) + \xi_1) + O(1)\sqrt{N}. \tag{16}$$

We analogically estimate $S(x^*(N),N)$ with few steps less and obtain

$$\widetilde{S}(x^*(N),N) = N(s'(x^*)+\xi_1) + O(1)N\epsilon(N).$$

Then we put above estimate together with (16) as a sums as in (17)

$$\frac{\sum_{i=1}^{\infty} e^{\frac{1}{N}\widetilde{S}'(\widetilde{x}^*(N),N)}i}{\sum_{i=1}^{\infty} e^{\frac{1}{N}S'(x^*(N),N)i}} = \frac{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i+O(1)(1/\sqrt{N})i}}{\sum_{i=1}^{\infty} e^{s'(x^*)i+O(1)\epsilon(N)i}} = \frac{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}(1+O(1)(1/\sqrt{N})i)}{\sum_{i=1}^{\infty} e^{s'(x^*)i}(1+O(1)\epsilon(N)i)},$$

and then we use Lemma 7 from the appendix and obtain the estimate

$$\frac{\sum_{i=1}^{\infty} e^{\frac{1}{N}\widetilde{S}'(\widetilde{x}^*(N),N)}i}{\sum_{i=1}^{\infty} e^{\frac{1}{N}S'(x^*(N),N)i}} = \frac{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}}{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}}\left(1+O(1)\frac{1}{\sqrt{N}}\right).$$

Finally we insert above estimate into (17) and obtain

$$M_{Y_N}(\xi) = \frac{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}}{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}}\left(1+O(1)\frac{1}{\sqrt{N}}\right)e^{\frac{1}{2}\hat{\xi}^T\hat{D}^2 s(x^*)^{-1}\hat{\xi}}\left(1+O\left(\frac{1}{h(N)^{1/2-3\delta}}\right)\right), \quad (17)$$

$$\text{when } \epsilon(N) \ll \frac{1}{h(N)^{1-3\delta}},$$

$$M_{Y_N}(\xi) = \frac{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}}{\sum_{i=1}^{\infty} e^{(s'(x^*)+\xi_1)i}}\left(1+O(1)\frac{1}{\sqrt{N}}\right)e^{\frac{1}{2}\hat{\xi}^T\hat{D}^2 s(x^*)^{-1}\hat{\xi}}\left(1+O\left(\sqrt{h(N)}\epsilon(N)\right)\right), \quad (18)$$

$$\text{when } \epsilon(N) \gg \frac{1}{h(N)^{1-3\delta}}, \quad (19)$$

and once we simplify estimate by including only the 'strongest' reminder term we get the final result. □

4 Partition function approximations

First we introduce a differentiable function $g : \Omega_N \to \mathbb{R}$ such that

$$\sup_{x\in\Omega_N} \|Dg(x)\| = g^{(1)} < \infty. \quad (20)$$

Lemma 2. *For defined above $S(x,N)$ with the maximum of type a)*

$$\left|\sum_{\Omega_N} g(x)e^{S(x,N)} - \frac{N^m}{d}g(x^*(N))e^{S(x^*(N),N)}\frac{(2\pi)^{m/2}}{\sqrt{|\det D^2S(x^*(N),N)|}}\right| \leq$$

$$\leq \frac{K}{N^{1/2-3\delta}}\frac{N^m}{d}|g(x^*(N))|e^{S(x^*(N),N)}\frac{(2\pi)^{m/2}}{\sqrt{|\det D^2S(x^*(N),N)|}},$$

vallid for $N > N_0$ where $\delta \in (0,\frac{1}{6})$ and d, K are some positive constants.

Proof. We start by decomposing the initial sum into two. One within the sphere $|x^*(N)-x| \leq \frac{1}{h(N)^{1/2-\delta}}$ and other one outside this sphere

$$\sum_{\Omega_N} g(x)e^{S(x,N)} = \sum_{\Omega'_N} g(x)e^{S(x,N)} + \sum_{\Omega''_N} g(x)e^{S(x,N)},$$

where Ω'_N represents the points in the domain inside the sphere and Ω''_N outside. Then for the first integral we expand function S with third order Taylor expansion at $x^*(N)$ and for the second integral we expand S with second order Taylor expansion and get

$$\begin{aligned}\sum_{\Omega_N} g(x)e^{S(x,N)} =& \sum_{\Omega'_N} g(x)e^{S(x^*(N),N)+(x-x^*)(N)D^2S(x^*(N),N)(x-x^*(N))+D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}}+\\ &+\sum_{\Omega''_N} g(x)e^{S(x^*(N),N)+(x-x^*(N))D^2S(x_{\theta_2(N)},N)(x-x^*(N))},\end{aligned}$$

where the first derivative vanish as $x^*(N)$ is a critical point of S. Then we expand the function g with first order Taylor approximation at $x^*(N)$

$$\begin{aligned}\sum_{\Omega_N} g(x)e^{S(x,N)} =& \sum_{\Omega'_N} g(x^*(N))e^{S(x^*(N),N)+(x-x^*)(N)D^2S(x^*(N),N)(x-x^*(N))+D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}}+\\ &+\sum_{\Omega'_N} g(x_{\theta_3(N)})^T(x-x^*N))e^{S(x*N,N)+(x-x^*)(N)D^2S(x^*(N),N)(x-x^*(N))}\times \\ & \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad (21)\\ &\times e^{D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}}+\sum_{\Omega''_N} g(x)e^{S(x^*(N),N)+(x-x^*(N))D^2S(x_{\theta_2(N)},N)(x-x^*(N))}. \\ & \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad (22)\end{aligned}$$

Since the third order term of the expansion of S consists of following expressions

$$\frac{\partial^3 S(x_{\theta_1(N)})}{\partial x_i \partial x_j \partial x_k}(x_i - x_i^*(N))(x_j - x_j^*(N))(x_k - x_k^*(N))$$

and by the equivalence of the norm in finite dimensional space we have for the max norm

$$|x_i - x_i^*(N)|K \leq \max_i |x_i - x_i^*(N)|K \leq |x - x^*(N)| \leq \frac{1}{h(N)^{1/2-\delta}}$$

hence

$$x_i - x^*(N) = \frac{\sigma}{h(N)^{1/2-\delta}}$$

where σ is some constant. Since $D^{\otimes 3}S$ by differentiating (2) is of order $h(N)$, this implies by increasing N expression in (4) can be made arbitraty small. Therefore we can expand the exponent with the third derivatives of S in (21)

$$e^{D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}} = 1 + O(D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}), \text{ as } N \to \infty,$$

and we substitute it in (21) and after some transformations we get

$$\begin{aligned}&\sum_{\Omega_N} g(x)e^{S(x,N)} - \sum_{\Omega'_N} g(x^*(N))e^{S(x^*(N),N)+(x-x^*)(N)D^2S(x^*(N),N)(x-x^*(N))} =\\ &\quad= \sum_{\Omega'_N} g(x^*(N))e^{S(x^*(N),N)+(x-x^*)(N)D^2S(x^*(N),N)(x-x^*(N))}O(D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3})+\\ &\quad+\sum_{\Omega'_N} Dg(x_{\theta_2(N)})^T(x-x^*N))e^{S(x^*(N),N)+(x-x^*)(N)D^2S(x^*(N),N)(x-x^*(N))+D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}}+\\ &\quad+\sum_{\Omega''_N} g(x)e^{S(x^*(N),N)+(x-x^*(N))D^2S(x_{\theta_2(N)},N)(x-x^*(N))}. \qquad (23)\end{aligned}$$

Now let us bound some expressions for above inequality.
Since we assumed (5), by the Shwarz inequality and properties of tensor product we have

$$\begin{aligned}&\left|D^{\otimes 3}S(x_{\theta_1(N)},N)(x-x^*(N))^{\otimes 3}\right| \leq \left\|D^{\otimes 3}S(x_{\theta_1(N)},N)\right\|\left|(x-x^*(N))^{\otimes 3}\right| \\ &\leq h(N)s^{(3)}|x-x^*(N)|^3 \leq s^{(3)}\frac{1}{h(N)^{1/2-3\delta}},\end{aligned} \tag{24}$$

where we used Lemma 1 and the last bound is due to fact that we consider $x \in \Omega'_N$, i.e. $|x-x^*(N)| \leq \frac{1}{h(N)^{1/2-\delta}}$.
Further, by (20) and Schwarz inequality

$$|Dg(x_{\theta_3(N)})^T(x-x^*N))| \leq g^{(1)}|x-x^*(N)| \leq g^{(1)}\frac{1}{N^{1/2-\delta}}. \tag{25}$$

where the last bound is since we consider $x \in \Omega'_N$.
Since D^2S is negative definite, by Schwartz inequality, assumption (4) and the inequality form appendix (80) we have

$$\begin{aligned}&(x-x^*(N))^TD^2S(x_{\theta_2(N)},N)(x-x^*(N)) = -|(x-x^*(N))^TD^2S(x_{\theta_2(N)},N)(x-x^*(N))| \leq \\ &\leq -\frac{1}{\|D^2S(x_{\theta_2(N)},N)^{-1}\|}|x-x^*(N)|^2 \leq -h(N)s'^{(2)}|x-x^*(N)|^2 < -h(N)^{2\delta}s'^{(2)},\end{aligned} \tag{26}$$

where last inequality due to fact that we take $x \in \Omega''_N$, i.e $|x-x^*(N)| > \frac{1}{h(N)^{1/2-\delta}}$.
Now we take absolute value on both sides of (23) and bound the RHS using (24), (25) and (26)

$$\begin{aligned}&\left|\sum_{\Omega_N} g(x)e^{S(x,N)} - \sum_{\Omega'_N} g(x^*(N))e^{S(x^*(N),N)+(x-x^*(N))^TD^2S(x^*(N),N)(x-x^*(N))}\right| \leq \\ &\quad\leq \sum_{\Omega'_N}|g(x^*(N))|e^{S(x^*(N),N)+(x-x^*(N))^TD^2S(x^*(N),N)(x-x^*(N))}Ks^{(3)}\frac{1}{h(N)^{1/2-3\delta}}+ \\ &\quad+\sum_{\Omega'_N} g^{(1)}\frac{1}{h(N)^{1/2-\delta}}e^{S(x^*(N),N)+(x-x^*(N))^TD^2S(x^*(N),N)(x-x^*(N))+s^{(3)}\frac{1}{h(N)^{1/2-3\delta}}}+ \\ &\quad+\sum_{\Omega''_N}\max_{x\in\Omega_N}|g(x)|e^{S(x^*(N),N)-h(N)^{2\delta}s'^{(2)}},\end{aligned}$$

and then we perform some rearrangements of expressions

$$\begin{aligned}&\left|\sum_{\Omega_N} g(x)e^{S(x,N)} - \sum_{\Omega'_N} g(x^*(N))e^{S(x^*(N),N)+(x-x^*(N))^TD^2S(x^*(N),N)(x-x^*(N))}\right| \leq \\ &\quad\leq \sum_{\Omega'_N}\frac{1}{h(N)^{1/2-3\delta}}|g(x^*(N))|e^{S(x^*(N),N)+(x-x^*(N))^TD^2S(x^*(N),N)(x-x^*(N))}\Bigg[Ks^{(3)}+ \\ &\quad+g^{(1)}\frac{1}{h(N)^{2\delta}}e^{s^{(3)}\frac{1}{h(N)^{1/2-3\delta}}}+h(N)^{1/2-3\delta}\frac{\max_{x\in\Omega_N}|g(x)|}{g(x^*(N))}\frac{\sum_{\Omega''_N}e^{-h(N)^{2\delta}s'^{(2)}}}{\sum_{\Omega'_N}e^{(x-x^*(N))^TD^2S(x^*(N),N)(x-x^*(N))}}\Bigg].\end{aligned} \tag{27}$$

Then we perform further bounding, but concerning only the last expression in the bracket of RHS. First we bound its denominator. We use the Schwartz inequality and assumption

(3)

$$\begin{aligned}\sum_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} &\geq \sum_{\Omega'_N} e^{-\|D^2 S(x^*(N),N)\||x-x^*(N)|^2} \geq \\ \geq \sum_{\Omega'_N} e^{-h(N)s^2|x-x^*(N)|^2} &\geq \sum_{\Omega'''_N} e^{-s^2} \geq e^{-s^{(2)}},\end{aligned} \tag{28}$$

where in the second to last step we reduced the domain of sumation such that $|x-x^*| \leq \frac{1}{h(N)^{1/2}}$ denoted as Ω'''_N which was possible since all the summands are positive. In the last step we reduced sum to one point.
Since the sum in the expression in the numerator does not depend on summed element, it is just number of elements in the sum. Therefore in can be bounded from above by the number of elements in the whole domain $\Omega_{N,E}$. As this number is rather complicated to calucate we further bound the domain by the $m-1$-dimensional hypercube of the edge size 1. Since the points are separated by the distance $1/N$ in any directions there are $(N+1)^{m-1}$ of such points. Hence we have

$$\sum_{\Omega''_N} 1 \leq (N+1)^{m-1}. \tag{29}$$

Now we substitute estimates (28) and (29) into (27) and further bound from above by fixing N in certain parts

$$\begin{aligned}&\left|\sum_{\Omega_N} g(x)e^{S(x,N)} - \sum_{\Omega'_N} g(x^*(N))e^{S(x^*(N),N)+(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}\right| \leq \\ &\quad\leq \sum_{\Omega'_N} \frac{1}{h(N)^{1/2-3\delta}}|g(x^*(N))|e^{S(x^*(N),N)+(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}\Bigg[Ks^{(3)}+ \\ &\quad+ g^{(1)}\frac{1}{h(N)^{2\delta}}e^{s^{(3)}\frac{1}{h(N_0)^{1/2-3\delta}}} + h(N_0)^{1/2-3\delta}(N_0+1)^{m-1}\frac{\max_{x\in\Omega_N}|g(x)|}{g(x^*(N))}e^{-h(N_0)^{2\delta}s'^{(2)}+s^{(2)}}\Bigg],\end{aligned}$$

and then denote expression in the bracket as K_0

$$\begin{aligned}&\left|\sum_{\Omega_N} g(x)e^{S(x,N)} - g(x^*(N))e^{S(x^*(N),N)}\sum_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}\right| \leq \\ &\quad\leq \frac{K_0}{h(N)^{1/2-3\delta}}|g(x^*(N))|e^{S(x^*(N),N)}\sum_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))},\end{aligned}$$

where the constant $\delta \in (0, 1/6)$ in order to estimate be valid.
The next step is to use the result of the Theorem 1 to estimate the sum with an integral. The volume of the grid size is equal to $1/N^m$ as it is a hypercube fo dimension m with edges $1/N$, therefore the parameter $d=1$. With appropriate calculations we get

$$\begin{aligned}&\left|\sum_{\Omega_N} g(x)e^{S(x,N)} - N^m g(x^*(N))e^{S(x^*(N),N)}\int_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}dx\right| \leq \\ &\quad\leq N^m\frac{h(N)}{N}\frac{K}{h(N)^{(1/2-\delta)(m+1)}}|g(x^*(N))|e^{S(x^*(N),N)}\left[1+\frac{K_0}{h(N)^{1/2-3\delta}}\right]+ \\ &\quad+ N^m\frac{K_0}{h(N)^{1/2-3\delta}}|g(x^*(N))|e^{S(x^*(N),N)}\int_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}.\end{aligned} \tag{30}$$

Further we represent the integral in above inequality as

$$
\begin{aligned}
&\int_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx = \\
&= \int_{\mathbb{R}^m} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx + \int_{\mathbb{R}^m/\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx,
\end{aligned}
$$

where the first integral we calculate using Lemma 1 from Analysis Appendix

$$
\begin{aligned}
&\int_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx = \\
&= \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}} + \int_{\mathbb{R}^m/\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx.
\end{aligned}
$$

We substitute above into (30) and together with some manipulations get

$$
\begin{aligned}
&\Big| \sum_{\Omega_N} g(x) e^{S(x,N)} - N^m g(x^*(N)) e^{S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N))|}} \Big| \leq \\
&\quad \leq N^m \frac{h(N)}{N} \frac{K}{h(N)^{(1/2-\delta)(m+1)}} |g(x^*(N))| e^{Nf(x^*(N),N)} \left[1 + \frac{K_0}{h(N)^{1/2-3\delta}} \right] + \\
&\quad + N^m |g(x^*(N))| e^{S(x^*(N),N)} \int_{\mathbb{R}^m/\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx + \\
&\quad + N^m \frac{K_0}{h(N)^{1/2-3\delta}} |g(x^*(N))| e^{S(x^*(N),N)} \left[\frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N))|}} + \right. \\
&\quad \left. + \int_{\mathbb{R}^m/\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} \right],
\end{aligned}
$$

and we bound remaining integral in the inequality using Lemma 5 setting from the Analysis Appendix $N = h(N)$ and $A = D^2 S/h(N)$

$$
\begin{aligned}
&\Big| \sum_{\Omega_N} g(x) e^{S(x,N)} - N^m g(x^*(N)) e^{S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N))|}} \Big| \leq \\
&\quad \leq N^m \frac{h(N)}{N} \frac{K}{h(N)^{(1/2-\delta)(m+1)}} |g(x^*(N))| e^{S(x^*(N),N)} \left[1 + \frac{K_0}{h(N)^{1/2-3\delta}} \right] + \\
&\quad + N^m e^{-h(N)^{2\delta}} |g(x^*(N))| e^{S(x^*(N),N)} \frac{1}{h(N)^{m/2} \sqrt{|\det D^2 S(x^*(N),N)|}} \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})} \times \\
&\quad \times \left[\frac{1}{h(N)} \left(\frac{1 + \sqrt{1 + 4h(N)^{-1+2\delta}}}{2} \right)^{m-1} + \frac{\Gamma(m)}{h(N)^m} \right] \left[1 + \frac{K_0}{h(N)^{1/2-3\delta}} \right] + \\
&\quad + N^m \frac{K_0}{h(N)^{1/2-3\delta}} |g(x^*(N))| e^{S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N))|}},
\end{aligned}
$$

and the last term is asymptotically 'strongest'. Let us bound the above expression by fixing $N = N_0$ in some expresions and introduce constants

$$\Big|\sum_{\Omega_N} g(x)e^{S(x,N)} - N^m g(x^*(N))e^{S(x^*(N),N)}\frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}}\Big| \leq$$

$$\leq \frac{1}{h(N)^{1/2-3\delta}} N^m |g(x^*(N))| e^{S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}} \Bigg\{ K_0 +$$

$$+ \frac{K}{h(N_0)^{(1/2-\delta)m+2\delta}} \frac{h(N_0)}{N_0} \frac{\sqrt{|\det D^2 S(x^*(N),N)|}}{(2\pi)^{m/2}} \left[1 + \frac{K_0}{h(N_0)^{1/2-3\delta}}\right] +$$

$$+ h(N_0)^{1/2-3\delta} e^{-h(N_0)^{2\delta}} \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})} \left[\frac{1}{h(N_0)} \left(\frac{1+\sqrt{1+4h(N_0)^{-1+2\delta}}}{2}\right)^{m-1} + \frac{\Gamma(m)}{h(N_0)^m}\right] \left[1 + \frac{K_0}{h(N_0)^{1/2-3\delta}}\right] \Bigg\}$$

Hence, finally we have

$$\Big|\sum_{\Omega_N} g(x)e^{S(x,N)} - N^m g(x^*(N))e^{S(x^*(N),N)}\frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}}\Big| \leq$$

$$\leq \frac{K'}{h(N)^{1/2-3\delta}} N^m |g(x^*(N))| e^{S(x^*(N),N)} \frac{(2\pi)^{m/2}}{\sqrt{|\det D^2 S(x^*(N),N)|}}$$

where $K' > 0$ is some constant. To ensure the validity of the estimate for we have a condition $\delta \in (0, 1/6)$. Therefore we get the result for the first case of the maximum.

□

Lemma 3. *For defined above $S(x,N)$ and the maximum of type b)*

$$\Big|\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \frac{N^{m-1}}{d} g(Tv^*(N))e^{S(Tv^*(N),N)} \sum_{i=0}^{\infty} e^{\frac{1}{N}S'(Tv^*(N),N)i} \frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2 S(Tv^*(N),N)|}}\Big| \leq$$

$$\leq \frac{K}{N^{1/2-3\delta}} \frac{N^{m-1}}{d} |g(Tv^*(N))| e^{S(Tv^*(N),N)} \sum_{i=0}^{\infty} e^{\frac{1}{N}S'(Tv^*(N),N)i} \frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2 S(Tv^*(N),N)|}}, \tag{31}$$

where $\delta \in (0, \min\{1/2m, 1/6\})$, d and K are some positive constants. The transformation $x = Tv$ is a rotation of the coordinate system such that the axis v_1 is orthogonal to the hyperplane of Ω_N on which maximum is attained. The matrix of derivatives $\hat{D}^2 f(Tv^*(N))$ is $m-1 \times m-1$ matrix along v_i, $i = 2, \ldots, m$ and s' is derivative of s along v_1. The estimate is valid for a subsequence of integers N which elements can be divided by some integer q, where $\frac{p}{q} = v_1^*$ with $v^* = T^{-1}x^*$. Further, we assume that $h(N) = N$ in the representation (2) of S.

Proof. We start with performing a rotation of the coordinate system $x = Tv$ in the way that axis of v_1 is orthogonal to the hyperplane which the maximum is attained and points of v_1 in the domain are increasing

$$\sum_{\Omega_N} g(x)e^{S(x,N)} = \sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)}.$$

Corresponding points of the domain of summation Ω_N are also rotated and we denote the rotated points of the domain as $T(\Omega_N)$. The maximum in the new coordinates will be denoted by $v^*(N)$. In the next step we decompose the sum into two. First one with points contained in the cylinder defined by two constraints. One along v_1, $|v_1 - v_1^*(N)| \le \frac{1}{h(N)^{1-\delta}}$ and the second along coordinates from $v_2, \dots, v_m$ denoted by $\hat{v}$, $|\hat{v} - \hat{v}^*(N)| \le \frac{1}{h(N)^{1/2-\delta}}$. This cylinder is denoted by $T(\Omega_N)'$. The second sum is simply over remaining points and we denote it by $T(\Omega_N)''$

$$\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} = \sum_{T(\Omega_N)'} g(Tv)e^{S(Tv,N)} + \sum_{T(\Omega_N)''} g(Tv)e^{S(Tv,N)}.$$

Then we expand function S into Taylor series, third order in the first sum and second in the second one

$$\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} = \sum_{T(\Omega_N)'} g(Tv)e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \times$$
$$\times\, e^{D^{\otimes 3} S(Tv_{\theta_1(N)},N)(v-v^*(N))^{\otimes 3}} +$$
$$+ \sum_{T(\Omega_N)''} g(Tv)e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv_{\theta_2(N)},N)(v-v^*(N))},$$

where the first the derivative S' is along v_1. Note that first derivatives along the coordinates $\hat{v}$ are equal to 0 since we expand in Taylor series at the maximum.
Next we expand into Taylor series of first order at $v^*(N)$ function g and get

$$\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} = \sum_{T(\Omega_N)'} g(Tv^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \times$$
$$\times\, e^{D^{\otimes 3} S(Tv_{\theta_1(N)},N)(v-v^*(N))^{\otimes 3}} +$$
$$+ \sum_{T(\Omega_N)'} Dg(Tv_{\theta_3(N)})^T (v - v^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 f(Tv^*(N),N)(v-v^*(N))} \times$$
$$\times\, e^{D^{\otimes 3} S(Tv_{\theta_1(N)},N)(v-v^*(N))^{\otimes 3}} +$$
$$+ \sum_{T(\Omega_{N,E})''} g(Tv)e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv_{\theta_2(N)},N)(v-v^*(N))}. \qquad (32)$$

Since the third order term of the expansion of S consists of following expressions

$$\frac{\partial^3 S(Tv_{\theta_1(N)})}{\partial v_i \partial v_j \partial v_k}(v_i - v_i^*(N))(v_j - v_j^*(N))(v_k - v_k^*(N)) \qquad (33)$$

and by the equivalence of the norm in finite dimensional space we have for the max norm

$$|v_i - v_i^*(N)|K \le \max_i |v_i - v_i^*(N)|K \le |\hat{v} - \hat{v}^*(N)| \le \frac{1}{h(N)^{1/2-\delta}}$$

hence

$$v_i - v_i^*(N) = \frac{\sigma}{h(N)^{1/2-\delta}} \qquad (34)$$

where σ is some constant. Since $D^{\otimes 3}S$ by differentiating (2) is of order $h(N)$. This implies that for $N > N_0$ expression in (33) can be made arbitrary small. Therefore we can expand the exponent with the third derivatives of S in (32)

$$e^{D^{\otimes 3} S(Tv_{\theta_1(N)})(v-v^*(N))^{\otimes 3}} = 1 + O\big(D^{\otimes 3} S(Tv_{\theta_1(N)})(v - v^*(N))^{\otimes 3}\big), \text{ as } N \to \infty, \qquad (35)$$

and we substitute it back to (32), and after some rearrangements

$$\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \sum_{T(\Omega_N)'} g(Tv^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} =$$

$$= \sum_{T(\Omega_N)'} g(Tv^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \times$$

$$\times O\big(D^{\otimes 3} S(Tv_{\theta_1(N)},N)(v-v^*(N))^{\otimes 3}\big) +$$

$$+ \sum_{T(\Omega_N)'} Dg(Tv_{\theta_3(N)})^T (v-v^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \times$$

$$\times \Big(1 + O\big(D^{\otimes 3} S(Tv_{\theta_1(N)},N)(v-v^*(N))^{\otimes 3}\big)\Big) +$$

$$+ \sum_{T(\Omega_N)''} g(Tv)e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv_{\theta_2(N)},N)(v-v^*(N))}. \tag{36}$$

Now let us bound some expressions for above inequality.
Since we assumed (5), by the Schwartz inequality, properties of tensor product and equivalence of norms we have

$$\big|D^{\otimes 3} S(Tv_{\theta_1(N)},N)(v-v^*(N))^{\otimes 3}\big| \le \big\|D^{\otimes 3} S(Tv_{\theta_1(N)})\big\| \big|(v-v^*(N))^{\otimes 3}\big| \le$$
$$\le h(N)s^{(3)}|v-v^*(N)|^3 \le s^{(3)}h(N)(\max_i |v_i - v_i^*|)^3 \le s^{(3)}\frac{1}{h(N)^{1/2-3\delta}}, \tag{37}$$

where the last bound is due to fact that we consider $v \in T(\Omega_N)'$, i.e. $|\hat{v} - \hat{v}^*(N)| \le \frac{1}{h(N)^{1/2-\delta}}$ and (34) is valid.
Further, by (20) and Schwartz inequality

$$|Dg(Tv_{\theta_3(N)})^T (v - v^* N))| \le g^{(1)}|v-v^*(N)| \le g^{(1)}\frac{1}{h(N)^{1/2-\delta}}. \tag{38}$$

where the last bound is since we consider $v \in T(\Omega_N)'$.
Since $D^2 S(Tv_{\theta_2(N)},N)$ is negative definite, by Schwarz inequality and assumption (4)

$$(v-v^*(N))^T D^2 S(Tv_{\theta_2(N)},N)(v-v^*(N)) = -|(v-v^*(N))^T D^2 S(Tv_{\theta_2(N)},N)(v-v^*(N))| \le$$
$$\le -\frac{1}{\|D^2 S(Tv_{\theta_2(N)},N)^{-1}\|}|v-v^*(N)|^2 < -h(N)^{2\delta}s'^{(2)}, \tag{39}$$

where last inequality due to fact that we take $v \in T(\Omega_N)''$, i.e $|v - v^*(N)| > \frac{1}{h(N)^{1/2-\delta}}$.
Next, we take absolute value on both sides of (36), apply triangle inequality on the RHS amd use estimates (37), (38) and (39)

$$\Bigg|\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \sum_{T(\Omega_N)'} g(Tv^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))}\Bigg|$$

$$\le \sum_{T(\Omega_N)'} |g(Tv^*(N))|e^{Nf(Tv^*(N),N)+Nf'(Tv^*(N),N)(v_1-v_1^*(N))+N(v-v^*(N))^T D^2 f(Tv^*(N),N)(v-v^*(N))} K s^{(3)}\frac{1}{h(N)^{1/2-3}}$$

$$+ \sum_{T(\Omega_N)'} g^{(1)}\frac{1}{h(N)^{1/2-\delta}}e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))+s^{(3)}\frac{1}{h(N)^{1/2-3\delta}}} +$$

$$+ \sum_{T(\Omega_N)''} |g(Tv)|e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))-h(N)^{2\delta}s'^{(2)}},$$

and then we perform some rearrangements of expressions

$$\begin{aligned}
&\Bigg| \sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \sum_{T(\Omega_N)'} g(Tv^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \Bigg| \\
&\quad \le \sum_{T(\Omega_N)'} \frac{1}{h(N)^{1/2-3\delta}} |g(Tv^*(N))| e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \times \\
&\quad \times \Bigg[K s^{(3)} + g^{(1)} \frac{1}{h(N)^{2\delta}} e^{s^{(3)} \frac{1}{h(N)^{1/2-3\delta}}} + h(N)^{1/2-3\delta} \frac{\max_{v\in T(\Omega_N)} |g(Tv)|}{g(Tv^*(N))} \times \\
&\quad \times \frac{\sum_{T(\Omega_N)''} e^{-h(N)^{2\delta} s'^{(2)}}}{\sum_{T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))}} \Bigg]. \qquad (40)
\end{aligned}$$

Then we perform further bounding, but concerning only the last expression in the bracket of RHS. First we bound its denominator. We use the Schwartz inequality and assumption (3)

$$\begin{aligned}
&\sum_{T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \ge \\
&\sum_{T(\Omega_N)'} e^{-|S'(Tv^*(N),N)||v_1-v_1^*(N)|-\|D^2 S(Tv^*(N),N)\||v-v^*(N)|^2} \ge \\
&\sum_{T(\Omega_N)'} e^{-|S'(Tv^*(N),N)||v_1-v_1^*(N)|-h(N)s^2|v-v^*(N)|^2} \ge \sum_{T(\Omega_N)'''} e^{-|S'(Tv^*(N),N)|/h(N)-s^2} \ge e^{-|S'(Tv^*(N),N)|/h(N)-F^{(2)}}, \qquad (41)
\end{aligned}$$

where in the second to last step we reduced the domain of summation such that $|v-v^*| \le \frac{1}{h(N)^{1/2}}$ and $|v_1 - v_1^*(N)| \le \frac{1}{h(N)}$ denoted as $T(\Omega_N)'''$ which was possible since all the summands are positive. In the last step we reduced sum to one point.

Since the sum in the expression in the numerator does not depend on summed element, it is just number of elements in the sum. Therefore in can be bounded from above by the number of elements in the whole domain $T(\Omega_N)$. As this number is rather complicated to calculate we further bound the domain by the m-dimensional hypercube of edge size 1. Since the points are separated by the distance $1/N$ in any directions there are $(N+1)^m$ of such points. Hence we have

$$\sum_{T(\Omega_N)''} 1 \le (N+1)^m. \qquad (42)$$

Now we substitute estimates (41) and (42) into (40) and further bound from above by fixing N

$$\begin{aligned}
&\Bigg| \sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \sum_{T(\Omega_N)'} g(Tv^*(N))e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \Bigg| \\
&\quad \le \sum_{T(\Omega_N)'} \frac{1}{h(N)^{1/2-3\delta}} |g(Tv^*(N))| e^{S(Tv^*(N),N)+S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \times \\
&\quad \Bigg[K s^{(3)} + g^{(1)} \frac{1}{h(N)^{2\delta}} e^{s^{(3)} \frac{1}{h(N_0)^{1/2-3\delta}}} + h(N_0)^{1/2-3\delta} (N_0+1)^m \frac{\max_{v\in T(\Omega_N)} |g(Tv)|}{g(Tv^*(N))} e^{-h(N_0)^{2\delta} s'^{(2)} + s^{(2)}} \Bigg],
\end{aligned}$$

and denoting expression in the bracket as K'

$$\left| \sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - g(Tv^*(N))e^{S(Tv^*(N),N)} \sum_{T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))} \right|$$
$$\leq \frac{K'}{h(N)^{1/2-3\delta}} |g(Tv^*(N))| e^{S(Tv^*(N),N)} \sum_{T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))+(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N))}, \tag{43}$$

where the constant $\delta \in (0, 1/6)$ in order to estimate be valid.
Then we perform further estimates. We start we following decomposition of the second expression in the exponent of the sums

$$(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N)) = \sum_{i,j=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_i \partial v_j}(v_i - v_i^*(N))(v_j - v_j^*(N)) =$$
$$= \sum_{i,j=2}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_i \partial v_j}(v_i - v_i^*(N))(v_j - v_j^*(N)) + \sum_{j=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_1 \partial v_j}(v_1 - v_1^*(N))(v_j - v_j^*(N)) +$$
$$+ \sum_{i=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_i \partial v_1}(v_i - v_i^*(N))(v_1 - v_1^*(N)) + \frac{\partial^2 S(Tv^*(N),N)}{\partial v_1 \partial v_1}(v_1 - v_1^*(N))(v_1 - v_1^*(N)).$$

Now as taking into account (34) and fact that $(v_1 - v_1^*(N)) = \frac{\sigma'}{h(N)^{1-\delta}}$ in the considered domain, we can estimate above

$$(v-v^*(N))^T D^2 S(Tv^*(N),N)(v-v^*(N)) = \sum_{i,j=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_i \partial v_j}(v_i - v_i^*(N))(v_j - v_j^*(N)) =$$
$$= (\hat{v} - \hat{v}^*(N))^T \hat{D}^2 S(Tv^*(N),N)(\hat{v} - \hat{v}^*(N)) +$$
$$+ \frac{\sigma'\sigma}{h(N)^{3/2-2\delta}} \left[\sum_{i=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_i \partial v_1} + \sum_{j=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_1 \partial v_j} \right] + \frac{\sigma'\sigma}{h(N)^{2-2\delta}} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_1 \partial v_1},$$

where $\hat{D}$ represents differentiation operator along vector (v_2, v_{m-1}). Since the second derivative of S represented by (2) is of order $h(N)$ the last two terms in the above estimate are tending to 0 as $N \to \infty$ and we can expand the exponent of this estimate in following way

$$e^{(\hat{v}-\hat{v}^*(N))^T \hat{D}^2 S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))} \times$$
$$\times\, e^{\frac{\sigma'\sigma}{h(N)^{3/2-2\delta}} \left[\sum_{i=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_i \partial v_1} + \sum_{j=1}^{m-1} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_1 \partial v_j} \right] + \frac{\sigma'\sigma}{h(N)^{2-2\delta}} \frac{\partial^2 S(Tv^*(N),N)}{\partial v_1 \partial v_1}} =$$
$$= e^{(\hat{v}-\hat{v}^*(N))^T \hat{D}^2 S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))} \left(1 + K_1 \frac{\sigma'\sigma}{h(N)^{1/2-2\delta}}\right) \left(1 + K_2 \frac{1}{h(N)^{1-2\delta}}\right) =$$
$$= e^{(\hat{v}-\hat{v}^*(N))^T \hat{D}^2 S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))} \left(1 + K_4 \frac{1}{h(N)^{1/2-2\delta}}\right),$$

where K_1 and K_2 are some constants and

$$K_4 = \sigma'\sigma + K_2 \frac{1}{h(N_0)^{1/2}} + \frac{K_1 \sigma' \sigma K_2}{h(N)^{1-2\delta}}.$$

Then we substitute above estimate into (43), perform some manipulations and obtain

$$\Bigg|\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - g(Tv^*(N))e^{S(Tv^*(N),N)}\sum_{T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))+(\hat{v}-\hat{v}^*(N))^T\hat{D}^2S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))}\Bigg|$$
$$\leq \frac{K''}{h(N)^{1/2-3\delta}}|g(Tv^*(N))|e^{S(Tv^*(N),N)}\sum_{T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))+(\hat{v}-\hat{v}^*(N))^T\hat{D}^2S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))}, \tag{44}$$

where

$$K'' = K' + K_4\frac{1}{h(N_0)^{\delta}} + K'K_4\frac{1}{h(N_0)^{1/2-2\delta}}.$$

Further, since the hyperplane on which is the maximum is orthogonal to axis v_1, all the points of the domain $T(\Omega_N)'$ can represented as a series of hyperplanes, instead of one lattice. Since the coeficients of the hyperplane on which maximum is are rational the points on each such hyperplane form a lattice and each such lattice is composed of the ponits of $T(\Omega_N)'$ which has a fixed coordinate of v_1. In the next step we reexpress the sum in the estimate (44)

$$\sum_{T(\Omega_N)'} = \sum_{v_1\in T(\Omega_N)'}\sum_{T(\Omega_N)'(v_1)}$$

which means that we first sum over coordinate v_1 and then for a fixed v_1 we sum over the points of $T(\Omega_N)'$.

We insert it into (44) and perform some rearreagements

$$\Bigg|\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - g(Tv^*(N))e^{S(Tv^*(N),N)}\sum_{v_1\in T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))}\times$$
$$\times\sum_{T(\Omega_N)'(v_1)} e^{(\hat{v}-\hat{v}^*(N))^T\hat{D}^2S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))}\Bigg| \leq$$
$$\leq \frac{K''}{h(N)^{1/2-3\delta}}|g(Tv^*(N))|e^{S(Tv^*(N),N)}\sum_{v_1\in T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))}\times$$
$$\times\sum_{T(\Omega_N)'(v_1)} e^{(\hat{v}-\hat{v}^*(N))^T\hat{D}^2S(Tv^*(N),N)(\hat{v}-\hat{v}^*(N))},$$

Now we perform approximation with the integral the last sum in the RHS and LHS. We do this step in the same way as for the previous case of maximum. The difference lies in in the dimension, its one less and with the structure of the grid, it might not be a regular hypercubes, therefore the parameter $d \neq 1$. After appropriate calculations we get

$$\Bigg|\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \frac{N^{m-1}}{d}g(Tv^*(N))e^{S(Tv^*(N),N)}\sum_{v_1\in T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))}\times$$
$$\times\frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2S(Tv^*(N),N)|}}\Bigg| \leq$$
$$\leq \frac{K'''}{h(N)^{1/2-3\delta}}\frac{N^{m-1}}{d}|g(Tv^*(N))|e^{S(Tv^*(N),N)}\sum_{v_1\in T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1-v_1^*(N))}\times$$
$$\times\frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2S(Tv^*(N),N)|}}, \tag{45}$$

where $K''' > 0$ and $d > 0$ are some constants. Note, that as previously, to ensure the validity of the estimate we have a condition $\delta \in (0, 1/6)$. Now, let us represent $v_1 = \frac{k+i}{N}$, $i = 0, \ldots, I_N$, where I_N corresponds to the last point within $T(\Omega_N)'$ and k is a positive integer such that $k = \lfloor N v_1^* \rfloor$. In this notation integer i represents the number of the point from the boundary where is the maximum v_1^*. Further we represent $v_1^* - v_1 = \frac{i}{N} + \varphi(N)$. The constant $\varphi(N) = \frac{N v_1^* - \lfloor N v_1^* \rfloor}{N}$ which is the distance form the maximum to the nearest point on the lattice on the axis v_1. Note that the constant $N\varphi(N)$ for oscillates on the interval $[0, 1]$ for various values of N. From the assumptions we have that v_1 is a rational, i.e. $v_1 = \frac{p}{q}$ for some integers p, q and q is a divisor of N hence $N v_1^* = \lfloor N v_1^* \rfloor$ hence the constant $\varphi(N) = 0$. Therefore we can represent the sum in (49) as

$$\sum_{v_1 \in T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1 - v_1^*(N))} = \sum_{i=0}^{I_N} e^{\frac{1}{N} S'(Tv^*(N),N) i}, \tag{46}$$

for a appropriate subsequence of N. Further, we can represnt the sum of above expression as a difference

$$\sum_{i=0}^{I_N} e^{\frac{1}{N} S'(Tv^*(N),N) i} = \sum_{i=0}^{\infty} e^{\frac{1}{N} S'(Tv^*(N),N) i} - \sum_{I_N}^{\infty} e^{\frac{1}{N} S'(Tv^*(N),N) i}, \tag{47}$$

Since I_N corresponds to biggest possible element v_i within $T(\Omega_N)'$, i.e. within $|v_1 - v_1^*| \leq N^{-1+\delta}$ hence elements larger than I_N corresponds to values of v_1 in the interval $|v_1 - v_1^*| > N^{-1+\delta}$ or equivalently $i > N^\delta$. Therefore one can estimate the second sum in the RHS of (4)

$$\begin{aligned} \sum_{I_N}^{\infty} e^{\frac{1}{N} S'(Tv^*(N),N) i} &= \sum_{i > N^\delta} e^{\frac{1}{N} S'(Tv^*(N),N) i} \leq \\ &\leq \int_{N^\delta}^{\infty} e^{\frac{1}{N} S'(Tv^*(N),N) v} dv + e^{\frac{1}{N} S'(Tv^*(N),N) N^\delta} - \lim_{v \to \infty} e^{\frac{1}{N} S'(Tv^*(N),N) v}, \end{aligned} \tag{48}$$

where the last estimate is due to standard approximation of sum with an integral.
Since S' due to representation (2) is of order $h(N)$ and we assumed that $h(N) = N$ the above estimate can be simplified and the integral calculated

$$= \frac{1}{s_1'(x) + \sigma_1(x)\epsilon(N)} e^{-|s_1'(x) + \sigma_1(x)\epsilon(N)| N^\delta} + e^{-|s_1'(x) + \sigma_1(x)\epsilon(N)| N^\delta}$$

where the S' expression in the exponent is negative since S' was negative.
Hence putting together above estimate with (46), (47) and (48) and adding absolute value appropriately one after some calculations gets

$$\left| \sum_{v_1 \in T(\Omega_N)'} e^{S'(Tv^*(N),N)(v_1 - v_1^*(N))} - \sum_{i=0}^{\infty} e^{\frac{1}{N} S'(Tv^*(N),N) i} \right| \leq K_1 e^{-|s_1'(x) + \sigma_1(x)\epsilon(N)| N^\delta}$$

where $K_1 = \sup_{x \in \Omega, N > N_0} \left(\frac{1}{s_1'(x) + \sigma_1(x)\epsilon(N)} + 1 \right)$.

Finally, we substitute above estimate into (49) and perform some calculations to obtain

$$\begin{aligned}&\Bigg|\sum_{T(\Omega_N)} g(Tv)e^{S(Tv,N)} - \frac{N^{m-1}}{d}g(Tv^*(N))e^{S(Tv^*(N),N)}\sum_{i=0}^{\infty} e^{\frac{1}{N}S'(Tv^*(N),N)i}\times\\ &\qquad\times\frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2S(Tv^*(N),N)|}}\Bigg| \leq\\ &\qquad\leq \frac{K_2}{h(N)^{1/2-3\delta}}\frac{N^{m-1}}{d}|g(Tv^*(N))|e^{S(Tv^*(N),N)}\sum_{i=0}^{\infty} e^{\frac{1}{N}S'(Tv^*(N),N)i}\times\\ &\qquad\times\frac{(2\pi)^{(m-1)/2}}{\sqrt{|\det D^2S(Tv^*(N),N)|}}, \end{aligned}\tag{49}$$

where constant K_2 is given by

$$K_2 = 1 + \frac{K_1 h(N_0)^{1/2-3\delta}}{K'''}e^{-|s_1'(x)+\sigma_1(x)\epsilon(N_0)|N_0^\delta} + K_1 e^{-|s_1'(x)+\sigma_1(x)\epsilon(N_0)|N_0^\delta}. \tag{50}$$

Hence we get the final result. □

5 Gaussian type integral approximation

Theorem 3. *For the function S with the maximum type a) on the domain Ω_N' containing the points $|x - x^*| \leq \frac{1}{h(N)^{1/2-\delta}}$ but with the distances between points given by $b_1/N, b_2/N, \dots, b_m/N$ for each direction in $\mathbb{R}^m$ we have following estimate*

$$\begin{aligned}&\left|\frac{N^m}{d}\int_{\Omega_N'} e^{(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))}dx - \sum_{x\in\Omega_N'} e^{(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))}\right| \leq\\ &\leq \frac{N^m}{d}\frac{h(N)}{N}\frac{K}{h(N)^{(1/2-\delta)(m+1)}},\end{aligned}$$

valid for $N > N_0$, where K is some positive constant, $\delta > 0$ is an arbitrary small constant and $d = \frac{1}{N}\prod_{i=1}^m b_i$ is a volume of the grid size.

Proof. We start by dividing the domain, i.e. sphere of the radius $\frac{1}{h(N)^{1/2-\delta}}$, into m-dimensional hypercuboid, with edges of size $\frac{b_1}{N}, \frac{b_2}{N}, \dots, \frac{b_m}{N}$. The hypercuboids are spread in the way that their middle points are the points of the lattice of the domain Ω_N. Since $x_i \in \{b_i, b_i/N, b_i/N, \dots, b_i(N-1)/N, 1\}$, $i = 1, \dots, m-1$ for Ω_N the considered hypercuboids fit that purpose. Note, that some hypercuboids on the edge of the domain might not contain the point from Ω_N' as their middle point is outside the domain but the have middle point within larger Ω_N. We write

$$\int_{\Omega_N'} e^{(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))}dx = \sum_{\omega\in\Omega_N'}\int_{\omega} e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}dx \tag{51}$$

where ω is a single hypercuboid and we have minus and absolute value since D^2S is negative definite for $x \in \Omega_N$ Now, for each summand we apply first order Taylor theorem at the middle point of the hypercube x_ω, i.e.

$$\begin{aligned}\int_\omega e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}dx = \int_\omega \Big\{ e^{-|(x_\omega-x^*(N))^T D^2 S(x^*(N),N)(x_\omega-x^*(N))|} + \\ + (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}} \Big\} dx,\end{aligned}$$

where $x_{\theta(\omega)}$ is between x_ω and x and D is a differential operator along x. Then we divide the integral into two and integrate one with constant function and obtain

$$\begin{aligned}\int_\omega e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}dx = \frac{d}{N^m} e^{-|(x_\omega-x^*(N))^T D^2 S(x^*(N),N)(x_\omega-x^*(N))|} + \\ + \int_\omega (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}} dx,\end{aligned}$$

and we put it back into (51) and obtain

$$\begin{aligned}\int_{\Omega'_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}dx = \sum_{\omega\in\Omega'_N} \frac{d}{N^m} e^{-|(x_\omega-x^*(N))^T D^2 S(x^*(N),N)(x_\omega-x^*(N))|} + \\ + \sum_{\omega\in\Omega'_N} \int_\omega (x_\omega - x)^T D e^{-N|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega')}} dx\end{aligned} \tag{52}$$

We divide the first sum over all hypercuboid in above equation into one over those which middle points points are in Ω'_N and those which middle points are outside Ω'_N and denote this as Ω''_N

$$\begin{aligned}\sum_{\omega\in\Omega'_{N,E}} \frac{1}{N^m} e^{-|(x_\omega-x^*(N))^T D^2 S(x^*(N),N)(x_\omega-x^*(N))|} = \\ = \frac{1}{N^m} \sum_{x\in\Omega'_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|} + \frac{1}{N^m} \sum_{x\in\Omega''_{N,}} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|},\end{aligned}$$

Including that we can rewrite (53) into

$$\begin{aligned}\int_{\Omega'_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}dx = \frac{d}{N^m} \sum_{x\in\Omega'_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|} + \\ + \sum_{\omega\in\Omega'_N} \int_\omega (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}} dx + \\ + \frac{d}{N^m} \sum_{x\in\Omega''_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}.\end{aligned} \tag{53}$$

Then we perform some rearrangements and add absolute value on the both sides, apply trinagle inequality

$$\left|\int_{\Omega'_N} e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}dx - \frac{d}{N^m}\sum_{x\in\Omega'_N} e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}\right| \le$$

$$\le \sum_{\omega'\in\Omega'_N}\left|\int_\omega (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}} dx\right| +$$

$$+\frac{d}{N^m}\sum_{x\in\Omega''_N} e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|} \tag{54}$$

Then we estimate each of the two sums on the RHS of the above inequality. Since ω is a hypercuboid the edge sizes $\frac{b_1}{N}, \frac{b_2}{N}, \ldots, \frac{b_m}{N}$, for any $x \in \omega$ we have that $|x_{i,\omega} - x_i| \le \frac{b_i}{N}$, $i = 1, \ldots, m-1$,and therefore by equivalence of max norm $|x_\omega - x| \le \max_i |x_{i,\omega} - x_i| \le \max_i \frac{b_i}{N}$. Hence we can bound the integral over the hypecuboids ω in following way

$$\int_\omega \left|(x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}} dx\Big|_{x=x_{\theta(\omega)}}\right| dx \le$$

$$\le \frac{bm}{N}\int_\omega \max_i \left|\frac{\partial}{\partial x_i} e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}}\right| dx, \tag{55}$$

where $\max_i b_i = b$ and we bounded the sum of derivatives by m maximal derivatives. Then we calculate the derivative

$$\frac{bm}{N}\int_\omega \max_i \left|\frac{\partial}{\partial x_i} e^{-|(x-x^*(N))^T D^2S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}}\right| dx =$$

$$= \frac{bm}{N}\int_\omega \max_i \left|\frac{\partial^2 S(x_{\theta(\omega)}, N)}{\partial x_i^2} 2(x_i - x_i^*(N)) + \sum_{j=1, j\neq i}^{m-1} \frac{\partial^2 S(x_{\theta(\omega)}, N)}{\partial x_i \partial x_j}(x_j - x_j^*(N))\right| \times$$

$$\times\, e^{-|(x_{\theta(\omega)} - x^*(N))^T D^2S(x^*(N),N)(x_{\theta(\omega)} - x^*(N))|} dx, \tag{56}$$

then we bound the expression under the integral, noting that $|x_i - x_i^*(N)| \le \frac{1}{h(N)^{1/2-\delta}}$ and calculate the integral itself

$$= \frac{bm}{N}\int_\omega \max_i \left|\frac{\partial^2 S(x_{\theta(\omega)}, N)}{\partial x_i^2} 2(x_i - x_i^*(N)) + \sum_{j=1, j\neq i}^{m-1} \frac{\partial^2 S(x_{\theta(\omega)}, N)}{\partial x_i \partial x_j}(x_j - x_j^*(N))\right| \times$$

$$\times\, e^{-|(x_{\theta(\omega)} - x^*(N))^T D^2S(x^*(N),N)(x_{\theta(\omega)} - x^*(N))|} dx \le$$

$$\le \frac{h(N)}{N}\frac{dbm}{N^m}\frac{K}{h(N)^{1/2-\delta}} e^{-|(x_{\theta(\omega)} - x^*(N))^T D^2S(x^*(N),N)(x_{\theta(\omega)} - x^*(N))|}, \tag{57}$$

where we used the fact that from the representation (2) $\frac{\partial^2 S}{\partial x^2}$ is of order $h(N)$.
In order to get rid of dependence of above estimate on the particular hypercube we introduce estimates. Since D^2S is negative definite, by Schwartz inequality, assumption (4) and the inequality form appendix (80) we have

$$(x_{\theta(\omega)} - x^*(N))^T D^2S(x_{\theta_2(N)}, N)(x_{\theta(\omega)} - x^*(N)) = -|(x_{\theta(\omega)} - x^*(N))^T D^2S(x_{\theta_2(N)}, N)(x_{\theta(\omega)} - x^*(N))| \le$$

$$\le -\frac{1}{\|D^2S(x_{\theta_2(N)}, N)^{-1}\|}|x_{\theta(\omega)} - x^*(N)|^2 \le 0$$

where last inequality due to fact that $|x_{\theta(\omega)} - x^*(N)| \geq 0$ for $x \in \Omega'_N$. Hence we can write

$$\begin{aligned}
&\frac{h(N)}{N}\frac{dbm}{N^m}\frac{K}{h(N)^{1/2-\delta}}e^{-|(x_{\theta(\omega)}-x^*(N))^T D^2 S(x^*(N),N)(x_{\theta(\omega)|}-x^*(N))} \leq \\
&\leq \frac{h(N)}{N}\frac{dbm}{N^m}\frac{K}{h(N)^{1/2-\delta}}. \qquad (58)
\end{aligned}$$

We combine the estimtes (55), (56), (57) and (58), then we put it into orginal sum over all hypercubes in (53) and obtain

$$\begin{aligned}
&\sum_{\omega \in \Omega'_N} \int_\omega (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|}\Big|_{x=x_{\theta(\omega)}} dx \leq \\
&\leq \frac{h(N)}{N}\frac{1}{N^m}\frac{K'}{h(N)^{1/2-\delta}} \sum_{\omega \in \Omega'_N} 1 \qquad (59)
\end{aligned}$$

Now we estimate the number of the hypercuboid ω. Note, that in the hypercube of the edge size $\frac{1}{h(N)^{1/2-\delta}}$ of dimension m there are contained $\prod_{i=1}^m \lfloor \frac{N}{b_i h(N)^{1/2-\delta}} \rfloor$ hypercuboids. The exact number of hypercuboid is determined by the position of the hypercube with the edge $\frac{1}{h(N)^{1/2-\delta}}$ on the grid of the hypercuboids. Some hypercuboids are contained partially. The total number of hypercuboids, those contained completely and partially can be bounded by adding extra hypercuboids. We add extra hypercuboid on each boundary and in each corner. Hence we have

$$\sum_\omega 1 \leq \prod_{i=1}^m \left(\lfloor \frac{N}{b_i h(N)^{1/2-\delta}} \rfloor + 2 \right) + 2^m,$$

where extra term inside the bracket corresponds to the boundary and the last term to the corners. We also can bound the $\prod_{i=1}^m \lfloor \frac{N}{b_i h(N)^{1/2-\delta}} \rfloor$ and obtain

$$\sum_\omega 1 \leq \left(\frac{N}{h(N)^{1/2-\delta} b} + 2 \right)^m + 2^m.$$

where $\max_i b_i = b$.
We combine above estimate with (59) and perform some simple calculations

$$\begin{aligned}
&\sum_{\omega \in \Omega'_N} \int_\omega (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}\Big|_{x=x_{\theta(\omega)}} dx \leq \\
&\leq \frac{h(N)}{N}\frac{K'}{h(N)^{(1/2-\delta)(m+1)}} \left[\left(\frac{1}{b} + \frac{2h(N)^{1/2-\delta}}{N} \right)^m + \left(\frac{2h(N)^{1/2-\delta}}{N} \right)^m \right],
\end{aligned}$$

Therefore, eventually we get

$$\sum_{\omega \in \Omega'_N} \int_\omega (x_\omega - x)^T D e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))}\Big|_{x=x_{\theta(\omega)}} dx \leq \frac{h(N)}{N}\frac{K''}{h(N)^{(1/2-\delta)(m+1)}}, \qquad (60)$$

valid for some $N > N_0$, where

$$K'' = K' \left[\left(\frac{1}{b} + \frac{2h(N_0)^{1/2-\delta}}{N_0} \right)^m + \left(\frac{2h(N_0)^{1/2-\delta}}{N_0} \right)^m \right].$$

For the second sum in the RHS of (54) we use the fact that the points $x \in \Omega''_N$ are such that $|x - x^*(N)| \geq \frac{1}{h(N)^{1/2-\delta}}$. Hence using Schwartz inequality and assumption (4) we get an estimate of the expression in the exponent

$$- |(x - x^*(N))^T D^2 S(x^*(N), N)(x - x^*(N)) \leq - \frac{1}{\|D^2 S(x^*(N), N)^{-1}\|}|x - x^*(N)|^2 \leq$$
$$\leq -h(N)^{2\delta} s'^{(2)},$$

and we have an estimate of whole expression

$$\sum_{x \in \Omega''_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|} \leq e^{-h(N)^{2\delta} s'^{(2)}} \sum_{x \in \Omega''_N} 1$$

Note that we consider only those $x \in \Omega''_N$ that are middle points of the hypercuboids which are partially contained in the domain Ω'_N. However we make a crude estimate and bound this number of hypercuboids by a number of hypercuboids which contains the hypercube of edge size 1, which contains whole domain Ω_N. This number is N^m. Therefore we have a final estimate of second sum

$$\sum_{x \in \Omega''_N} e^{-|(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))|} \leq e^{-h(N)^{2\delta} s'^{(2)}} N^m. \tag{61}$$

Finally, we substitute estimates (60) and (61) into (54)

$$\left| \int_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx - \frac{d}{N^m} \sum_{x \in \Omega'_N} e^{(x-x^*(N))^T D^2 f(x^*(N),N)(x-x^*(N))} \right| \leq$$
$$\leq \frac{h(N)}{N} \frac{K''}{h(N)^{(1/2-\delta)(m+1)}} + e^{-h(N)^{2\delta} s'^{(2)}} N^m.$$

We multiply whole expression by $\frac{N^m}{d}$ and perform some manipulations

$$\left| \frac{N^m}{d} \int_{\Omega'_N} e^{(x-x^*(N))^T D^2 S(x^*(N),N)(x-x^*(N))} dx - \sum_{x \in \Omega'_N} e^{(x-x^*(N))^T D^2 f(x^*(N),N)(x-x^*(N))} \right| \leq$$
$$\leq \frac{N^m}{d} \frac{h(N)}{N} \frac{K'''}{h(N)^{(1/2-\delta)(m+1)}}.$$

where K''' is given by

$$K''' = 1 + e^{-h(N_0)^{2\delta} s'^{(2)}} \frac{N_0}{h(N_0)} \frac{h(N_0)^{(1/2-\delta)(m+1)}}{K'''},$$

valid for $N > N_0$.

□

6 Additional estimates

Preposition 1. *For the function* S *with maximum of typa a) and function* $\widetilde{S}(x, N) = S(x, N) + \sqrt{h(N)}(x^* - x)^T \xi$ *which has maximum at the point* $x^*(N)$ *we have following* *results*

$$\widetilde{x}^*(N) - x^*(N) = D^2 s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\bigg(\frac{1}{h(N)}\bigg), \tag{62}$$

$$\begin{aligned}\widetilde{S}(\widetilde{x}^*(N), N) &= S(x^*(N), N) - \frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi + O(1)\frac{1}{\sqrt{h(N)}} && \textit{when } \epsilon(N) \ll \frac{1}{h(N)},\\ \widetilde{S}(\widetilde{x}^*(N), N) &= S(x^*(N), N) - \frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi + O(1)\sqrt{h(N)}\epsilon(N) && \textit{when } \epsilon(N) \gg \frac{1}{h(N)},\end{aligned} \tag{63}$$

$$\frac{\sqrt{\det D^2 S(x^*(N), N)}}{\sqrt{\det D^2 \widetilde{S}(\widetilde{x}^*(N), N)}} = 1 + O\bigg(\frac{1}{\sqrt{h(N)}}\bigg). \tag{64}$$

as $N \to \infty$*, where for all results we assumed that* $\epsilon(N) \ll \frac{1}{\sqrt{h(N)}}$.

Proof. We begin with the proof of the first estimate.
Let us take first order Taylors expansion of the vector function $D\widetilde{S}(\widetilde{x}^*(N), N)$ at the point $x^*(N)$

$$D\widetilde{S}(\widetilde{x}^*(N), N) = DS(x^*(N), N) - \sqrt{h(N)}\xi + D^2 S(x_\theta(N), N)^T(\widetilde{x}^*(N) - x^*(N)).$$

where $x_\theta(N)$ is somewhere between $\widetilde{x}^*(N)$ and $x^*(N)$.
Since $x^*(N)$ and $\widetilde{x}^*(N)$ are a critical points, of respectively $S(x, N)$ and $\widetilde{S}(x, N)$ hence $DS(x^*(N), N) = D\widetilde{S}(\widetilde{x}^*(N), N) = o$, where $x = o$ is a vector of origin. Furthermore as $S \in C^2$ the order of differentiation does not matter we have that $D^2 S(x_\theta(N), N)^T = D^2 S(x_\theta(N), N)$. Hence following equality is valid

$$D^2 f_l(x_\theta(N), N)(\widetilde{x}^*(N) - x^*(N)) = \sqrt{h(N)}\xi.$$

Then we take define a lower bound as in (80) and have a bound due to (4) and after small transformation obtain

$$\big|\widetilde{x}^*(N) - x^*(N)\big| \leq \frac{(s'^{(2)})^{-1}|\xi|}{\sqrt{h(N)}} \tag{65}$$

Note that for different value of ξ the point $x_\theta(N)$ might be altered, but this does not influence the constant $s'^{(2)}$ as it is defined as a supremum over whole domain.
Further, we use second order Taylors expansion of $D\widetilde{S}(\widetilde{x}^*(N), N)$ at $x^*(N)$

$$\begin{aligned}D\widetilde{S}(\widetilde{x}^*(N), N) =& DS(x^*(N), N) - \sqrt{h(N)}\xi + D^2 S(x^*(N), N)(\widetilde{x}^*(N) - x^*(N)) + \\ &+ \frac{1}{2}D^3 S(x_\theta(N))(\widetilde{x}^*(N) - x^*(N))^{\otimes 2}\end{aligned}$$

where $x_\theta(N)$ is some point between $\widetilde{x}^*(N)$ and $x^*(N)$. Again, we have that $D\widetilde{S}(\widetilde{x}^*(N), N) = DS(x^*(N), N) = o$.
Now we transform above expansion and get the upper bound as in (82) bound it as in (5) and with some calculations obtain

$$\Big|D^2 S(x^*(N), N)(\widetilde{x}^*(N) - x^*(N)) - \sqrt{h(N)}\xi\Big| \leq h(N)\frac{s^{(3)}}{2}\big|\widetilde{x}^*(N) - x^*(N)\big|^2,$$

and then substitute estimate (65) in the RHS and obtain

$$\left|D^2S(x^*(N),N)(\widetilde{x}^*(N)-x^*(N))-\sqrt{h(N)}\xi\right|\leq\frac{s^{(3)}(s'^{(2)})^{-2}|\xi|}{2}.$$

Next we transform LHS of above inequality, bound the second derivative matrix and as a result obtain a lower bound of above

$$\left|(\widetilde{x}^*(N)-x^*(N))-D^2S(x^*(N),N)^{-1}\sqrt{h(N)}\xi\right|\leq\frac{s^{(3)}(s'^{(2)})^{-3}|\xi|}{2h(N)}. \tag{66}$$

Next we apply Taylor's theorem of the first order at x^* to $D^2S(x^*(N),N)^{-1}$ we get

$$D^2S(x^*(N),N)^{-1}=D^2S(x^*,N)^{-1}+D^2S(x_\theta(N),N)^{-1}(x^*(N)-x^*).$$

and then we transform it apply absolute values on both side and trinagle inequality

$$|D^2S(x^*(N),N)^{-1}-D^2S(x^*,N)^{-1}|\leq\|D^2S(x_\theta(N),N)^{-1}\||x^*(N)-x^*|.$$

and again transform

$$\frac{1}{h(N)}\|D^2S(x_\theta(N),N)^{-1}\|^{-1}|D^2S(x^*(N),N)^{-1}-D^2S(x^*,N)^{-1}|\leq\frac{1}{h(N)}|x^*(N)-x^*|.$$

Finaly we use the property (4) and (6) and obtain

$$|D^2S(x^*(N),N)^{-1}-D^2S(x^*,N)^{-1}|\leq\frac{1}{h(N)}(s'^{(2)})^{-1}O(1)\epsilon(N). \tag{67}$$

Now we perform some transformations on $D^2S(x^*,N)^{-1}$ using (7)

$$D^2S(x^*,N)^{-1}=\frac{1}{h(N)}\big[D^2s(x^*)+\sigma(x^*)\epsilon(N)\big]^{-1}=\frac{1}{h(N)}D^2s(x^*)^{-1}\big[I+\sigma(x^*)\epsilon(N)D^2s(x^*)\big]^{-1}$$

where I is $m\otimes m$ identity matrix. Now we apply the asymptotic version of second result of the Theorem 5 from the Appendix

$$(I+\epsilon A)^{-1}=I+O(\epsilon),\ \epsilon\to 0,$$

Hence we have that

$$D^2S(x^*,N)^{-1}=\frac{1}{h(N)}D^2s(x^*)^{-1}+\frac{1}{h(N)}D^2s(x^*)^{-1}\sigma(x^*)O(\epsilon(N))D^2s(x^*)$$

which can be simplified to

$$D^2S(x^*,N)^{-1}=\frac{1}{h(N)}D^2s(x^*)^{-1}+\frac{1}{h(N)}O(\epsilon(N))$$

since all the constants are contained in big 'O'. Then we substitute above estimate into (67) and after some calculations get

$$|D^2S(x^*(N),N)^{-1}-\frac{1}{h(N)}D^2s(x^*)^{-1}|\leq\frac{\epsilon(N)}{h(N)}O(1). \tag{68}$$

and finally we substitute above into (66) and after some manipulations obtain

$$\left|(\widetilde{x}^*(N)-x^*(N))-\frac{1}{\sqrt{h(N)}}D^2s(x^*)^{-1}\xi\right|\le\frac{s^{(3)}(s'^{(2)})^{-3}|\xi|}{2h(N)}+\frac{\epsilon(N)|\xi|}{\sqrt{h(N)}}O(1).$$

Now, since we assumed $\epsilon(N)\ll\frac{1}{\sqrt{h(N)}}$ we have

$$\left|(\widetilde{x}^*(N)-x^*(N))-\frac{1}{\sqrt{h(N)}}D^2s(x^*)^{-1}\xi\right|\le K\frac{1}{h(N)},$$

where $K=\frac{s^{(3)}(s'^{(2)})^{-3}|\xi|}{2}+\epsilon(N_0)\sqrt{h(N_0)}|\xi|O(1)$.
Now we prove the second result of this preposition.
We start by substituting the formula for $\widetilde{x}^*(N)$ given by the first result of the preposition into the explicit expression for $\widetilde{S}(\widetilde{x}^*(N),N)$ given in the statement of the preposition

$$\begin{aligned}\widetilde{S}(\widetilde{x}^*(N),N)&=S\left(x^*(N)+D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right),N\right)-\\&-\sqrt{h(N)}\xi^T\left(x^*(N)+D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)-x^*\right).\end{aligned}\tag{69}$$

Then expand the function S in RHS using 3-rd order Taylor expansion at the point $x^*(N)$

$$\begin{aligned}&S\left(x^*(N)+D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right),N\right)=\\&=S(x^*(N),N)+DS(x^*(N),N)\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)+\\&+\frac{1}{2}\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)^TD^2S(x^*(N),N)\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)+\\&+\frac{1}{6}D^{\otimes 3}S(x_{\theta_1}(N),N)\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)^{\otimes 3},\end{aligned}$$

where $x_{\theta_1}(N)$ is some point between $\widetilde{x}^*(N)$ and $x^*(N)$. Further, we substitute above in the previous equality (69) an note that $DS(x^*(N),N)=o$

$$\begin{aligned}&\widetilde{S}(\widetilde{x}^*(N),N)=S(x^*(N),N)-\sqrt{h(N)}\xi^T\left(x^*(N)+D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)-x^*\right)+\\&+\frac{1}{2}\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)^TD^2S(x^*(N),N)\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)+\\&+\frac{1}{6}D^{\otimes 3}S(x_{\theta_1}(N),N)\left(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}}+O\left(\frac{1}{h(N)}\right)\right)^{\otimes 3}.\end{aligned}$$

Next, we find the upper bound as in (78) and additionally bound the constant due to (5)

$$\Big|\widetilde{S}(\widetilde{x}^*(N),N) - S(x^*(N),N) + \sqrt{h(N)}\xi^T\Big(x^*(N) + D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big) - x^*\Big) -$$

$$- \frac{1}{2}\Big(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big)^T D^2S(x^*(N),N)\Big(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big)\Big| \le \tag{70}$$

$$\le \frac{1}{6}h(N)s^{(3)}\Big|D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big|^3,$$

Now use Taylors Theorem on $D^2S(x^*(N),N)$ at x^*

$$D^2S(x^*(N),N) = D^2S(x^*,N) + D^{\otimes 3}S(x_\theta(N),N)(x^*(N) - x^*),$$

and then we transform above estimate and bound the third derivative using (79) and (5)

$$|D^2S(x^*(N),N) - D^2S(x^*,N)| \le h(N)s^{(3)}|x^*(N) - x^*|,$$

and finally we use the estimate (6) and obtain

$$D^2S(x^*(N),N) = D^2S(x^*,N) + O(1)h(N)\epsilon(N), \tag{71}$$

where the big 'O' contains the constant $s^{(3)}$. Next we by differentiation of the representation (7) obtain other form of $D^2S(x^*,N)$

$$D^2S(x^*,N) = h(N)[D^2s(x^*) + D\sigma(x^*)\epsilon(N)],\ \ N \to \infty$$

and substitute it into (71)

$$D^2S(x^*(N),N) = h(N)D^2s(x^*) + O(1)h(N)\epsilon(N),$$

where again we included constant in big 'O'. Finally we and we substitute it into (70) together with the result (6) and perform some basic calculations

$$\Big|\widetilde{S}(\widetilde{x}^*(N),N) - S(x^*(N),N) + \xi^T D^2s(x^*)^{-1}\xi + \frac{1}{\sqrt{h(N)}}\xi^T O(1) + \sqrt{h(N)}\epsilon(N)\xi^T O(1) -$$

$$- \frac{1}{2}\Big(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big)^T h(N)D^2s(x^*(N))\Big(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big) -$$

$$- \frac{1}{2}\Big(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big)^T h(N)O(1)\epsilon(N)\Big(D^2s(x^*)^{-1}\frac{\xi}{\sqrt{h(N)}} + O\Big(\frac{1}{h(N)}\Big)\Big)\Big| \le$$

$$\le \frac{1}{6}\frac{s^{(3)}}{\sqrt{h(N)}}\Big|D^2s(x^*)^{-1}\xi + O\Big(\frac{1}{\sqrt{h(N_0)}}\Big)\Big|^3,$$

and perform further transformations, such as triangle and Schwarz inequalities and use of (82) and others

$$\Big|\widetilde{S}(\widetilde{x}^*(N),N) - S(x^*(N),N) + \frac{1}{2}\xi^T D^2s(x^*)^{-1}\xi\Big| \le$$

$$\le \frac{1}{6}\frac{s^{(3)}}{\sqrt{h(N)}}\Big|D^2s(x^*)^{-1}\xi + O\Big(\frac{1}{\sqrt{h(N)}}\Big)\Big|^3 + \epsilon(N)|O(1)|\frac{1}{2}\Big|D^2s(x^*)^{-1}\xi + O\Big(\frac{1}{\sqrt{N}}\Big)\Big|^2 +$$

$$+ \frac{1}{\sqrt{h(N)}}|\xi||O(1)| + \sqrt{h(N)}\epsilon(N)|\xi^T O(1)| + \frac{1}{\sqrt{h(N)}}|O(1)||\xi| + \frac{1}{2h(N)}|O(1)||D^2s(x^*)|.$$

Since we have restriction $\epsilon(N) \ll \frac{1}{\sqrt{h(N)}}$ we can reargange above inequality into

$$\left|\widetilde{S}(\widetilde{x}^*(N), N) - S(x^*(N), N) + \frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi\right| \leq$$
$$\leq \frac{1}{\sqrt{h(N)}}\left[\frac{1}{6}s^{(3)}\left|D^2 s(x^*)^{-1}\xi + O\left(\frac{1}{\sqrt{h(N)}}\right)\right|^3 + \sqrt{h(N)}\epsilon(N)|O(1)|\frac{1}{2}\left|D^2 s(x^*)^{-1}\xi + O\left(\frac{1}{\sqrt{N}}\right)\right|^2 +\right.$$
$$\left.+ |\xi||O(1)| + |O(1)||\xi| + \frac{1}{2\sqrt{h(N)}}|O(1)||D^2 s(x^*)|\right] + \sqrt{h(N)}\epsilon(N)|\xi^T O(1)|.$$

and other cases of $\epsilon(N)$ would result in the situation that second error therm would be infinite lub just non zero. Further within the case $\epsilon(N) \ll \frac{1}{\sqrt{h(N)}}$ the above inequality will have two subcases

$$\left|\widetilde{S}(\widetilde{x}^*(N), N) - S(x^*(N), N) + \frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi\right| \leq \frac{K_1}{\sqrt{h(N)}} \qquad \text{when } \epsilon(N) \ll \frac{1}{h(N)},$$
$$\left|\widetilde{S}(\widetilde{x}^*(N), N) - S(x^*(N), N) + \frac{1}{2}\xi^T D^2 s(x^*)^{-1}\xi\right| \leq K_2\sqrt{h(N)}\epsilon(N) \quad \text{when } \epsilon(N) \gg \frac{1}{h(N)},$$

with

$$K_1 = \frac{1}{6}s^{(3)}\left|D^2 s(x^*)^{-1}\xi + O\left(\frac{1}{\sqrt{h(N_0)}}\right)\right|^3 + \sqrt{h(N_0)}\epsilon(N_0)|O(1)|\frac{1}{2}\left|D^2 s(x^*)^{-1}\xi + O\left(\frac{1}{\sqrt{N_0}}\right)\right|^2 +$$
$$+ |\xi||O(1)| + |O(1)||\xi| + \frac{1}{2\sqrt{h(N_0)}}|O(1)||D^2 s(x^*)| + h(N_0)\epsilon(N_0)|\xi^T O(1)|,$$
$$K_2 = \frac{1}{h(N)\epsilon(N)}\left[\frac{1}{6}s^{(3)}\left|D^2 s(x^*)^{-1}\xi + O\left(\frac{1}{\sqrt{h(N)}}\right)\right|^3 + \sqrt{h(N)}\epsilon(N)|O(1)|\frac{1}{2}\left|D^2 s(x^*)^{-1}\xi + \frac{O(1)}{\sqrt{N}}\right|^2 +\right.$$
$$\left.+ |\xi||O(1)| + |O(1)||\xi| + \frac{1}{2\sqrt{h(N)}}|O(1)||D^2 s(x^*)|\right] + |\xi^T O(1)|,$$

hence we get the second result.
Finally, we now prove the Prepositions third result. First we take first order Taylor approximation of $D^2\widetilde{S}(x^*(N), N)$ at $\widetilde{x}^*(N)$

$$D^2\widetilde{S}(x^*(N), N) = D^2\widetilde{S}(\widetilde{x}^*(N), N) + D^{\otimes 3}S(x_\theta(N), N)(\widetilde{x}^*(N) - x^*(N), N).$$

Since $D^2\widetilde{S}(\widetilde{x}^*(N), N) = D^2 S(\widetilde{x}^*(N), N)$ as the term $\sqrt{h(N)}(x^* - x)^T\xi$ vanish in the second derivatives of x. Then we bound it as in (82) and (5)

$$\left|D^2 S(x^*(N), N) - D^2\widetilde{S}(\widetilde{x}^*(N), N)\right| \leq h(N)s^{(3)}|\widetilde{x}^*(N) - x^*(N)|.$$

Next, we substitute the first estimate of this preposition, i.e. for $\widetilde{x}^*(N) - x^*(N)$

$$\left|D^2 S(x^*(N), N) - D^2\widetilde{S}(\widetilde{x}^*(N), N)\right| \leq \sqrt{h(N)}s^{(3)}|D^2 s(x^*)^{-1}\xi + O(1/\sqrt{h(N)})|.$$

Now we divide inequality by $|D^2\widetilde{S}(\widetilde{x}^*(N), N)|$ and obtain

$$\left|\frac{D^2 S(x^*(N), N)}{D^2\widetilde{S}(\widetilde{x}^*(N), N)} - I\right| \leq \sqrt{h(N)}\frac{s^{(3)}|D^2 f_l(x^*)^{-1}\xi + O(1/\sqrt{h(N)})|}{|D^2\widetilde{S}(\widetilde{x}^*(N), N)|}. \tag{72}$$

Now we apply Taylors theorem of the first order at x^* for the function in the denominator of RHS

$$D^2\widetilde{S}(\widetilde{x}^*(N),N) = D^2S(x^*,N) + D^2S(x_\theta(N),N)(\widetilde{x}^*(N)-x^*)$$

Then we transform above equality into

$$|D^2\widetilde{S}(\widetilde{x}^*(N),N) - D^2S(x^*,N)| \le \|D^2S(x_\theta(N),N)\|(|\widetilde{x}^*(N)-x^*(N)|+|x^*(N)-x^*|),$$

and we bound the RHS using the first result of the Preposition and (3), (6)

$$|D^2\widetilde{S}(\widetilde{x}^*(N),N)-D^2S(x^*,N)| \le \sqrt{h(N)}\big[D^2s(x^*)^{-1}\xi+O(1/\sqrt{h(N)})+\sqrt{h(N)}\epsilon(N)|O(1)|\big].$$

Next we differentiate the representation (7), substitute it into above inequality and after some calculations obtain

$$|D^2\widetilde{S}(\widetilde{x}^*(N),N)-h(N)D^2s(x^*)| \le \sqrt{h(N)}\big[D^2s(x^*)^{-1}\xi+O(1/\sqrt{h(N)})+\sqrt{h(N)}\epsilon(N)(|O(1)|+\sigma(x^*))\big].$$

and finally due to fact $\epsilon(N) \ll 1/\sqrt{h(N)}$ we can get an estimate

$$|D^2\widetilde{S}(\widetilde{x}^*(N),N)-h(N)D^2s(x^*)| \le \sqrt{h(N)}\big[D^2s(x^*)^{-1}\xi+O(1/\sqrt{h(N_0)})+\sqrt{h(N_0)}\epsilon(N_0)(|O(1)|+\sigma(x^*))\big].$$

therefore we have following estimate after appropriately applying the triangle inequality

$$|D^2\widetilde{S}(\widetilde{x}^*(N),N)| \ge h(N)|D^2s(x^*)| - K\sqrt{h(N)},$$

where K_1 is some positive constant. We substitute above into (72)

$$\left|\frac{D^2S(x^*(N),N)}{D^2\widetilde{S}(\widetilde{x}^*(N),N)} - I\right| \le \frac{1}{\sqrt{h(N)}}\frac{s^{(3)}|D^2f_l(x^*)^{-1}\xi+O(1/\sqrt{h(N)})|}{|D^2s(x^*)|-K/\sqrt{h(N)}}.$$

hence the RHS can be bounded for some $N > N_0$

$$\frac{1}{\sqrt{h(N)}}\frac{s^{(3)}|D^2f_l(x^*)^{-1}\xi+O(1/\sqrt{h(N)})|}{|D^2s(x^*)|-K/\sqrt{h(N)}} \le \frac{1}{\sqrt{h(N)}}\frac{s^{(3)}|D^2f_l(x^*)^{-1}\xi+O(1/\sqrt{h(N_0)})|}{|D^2s(x^*)|-K/\sqrt{h(N_0)}}$$

therefore we get that

$$\left|\frac{D^2S(x^*(N),N)}{D^2\widetilde{S}(\widetilde{x}^*(N),N)} - I\right| \le \frac{K'}{\sqrt{h(N)}}.$$

In a different form we can write above as

$$\frac{D^2S(x^*(N),N)}{D^2\widetilde{S}(\widetilde{x}^*(N),N)} = I + \frac{\sigma}{\sqrt{N}}. \tag{73}$$

where $|\sigma| \le K'$.
Now we take following

$$\left|\frac{\sqrt{\det D^2S(x^*(N),N)}}{\sqrt{\det D^2\widetilde{S}(\widetilde{x}^*(N),N)}} - 1\right| = \left|\sqrt{\det\left(\frac{D^2S(x^*(N),N)}{D^2\widetilde{S}(\widetilde{x}^*(N),N)}\right)} - 1\right|, \tag{74}$$

which is valid by the properties of the determinant.
Further, the first order Taylor expansion of the function $f(x) = \sqrt{x}$ at 1 is

$$\sqrt{x} = 1 + (a)^{-1/2}(x-1)$$

where $a \in (x, 1)$. We apply above expansion to (74) with

$$x = \det\left(\frac{D^2S(x^*(N), N)}{D^2\widetilde{S}(\widetilde{x}^*(N), N)}\right),$$

and obtain

$$\left|\sqrt{\det\left(\frac{D^2S(x^*(N), N)}{D^2\widetilde{S}(\widetilde{x}^*(N), N)}\right)} - 1\right| = \frac{1}{\sqrt{a}}\left|\det\left(\frac{D^2S(x^*(N), N)}{D^2\widetilde{S}(\widetilde{x}^*(N), N)}\right) - 1\right|, \tag{75}$$

where $a \in (\min(x, 1), \max(x, 1))$.
From Theorem 4 from Appendix we have that

$$\left|\det\left(I + \frac{\sigma}{\sqrt{N}}\right) - 1 - \frac{\operatorname{tr}(\sigma)}{\sqrt{N}}\right| \leq \frac{K''}{N}$$

for some positive K''. This can be transformed into

$$\left|\det\left(I + \frac{\sigma}{\sqrt{N}}\right) - 1\right| \leq \frac{K'''}{\sqrt{N}},$$

with

$$K''' = |\operatorname{tr}(\sigma)| + \frac{K_3}{N_0}.$$

Hence, taking into account (73) and combining (74), (75) with above inequality one gets

$$\left|\frac{\sqrt{\det D^2S(x^*(N), N)}}{\sqrt{\det D^2\widetilde{S}(\widetilde{x}^*(N), N)}} - 1\right| \leq \frac{K'''}{\sqrt{h(N)a}}$$

which is the third result. □

A Analysis

Theorem 4 (Taylor's expansion)**.** *Suppose that f is a real function on the nontrivial convex closed set $A \in \mathbb{R}^m$, n is a positive integer, $f(n-1)$ is continuous on A, $f^{(n)}(t)$ exists for every $t \in A$. Then there exists a point x_θ between x^* and x, such that*

$$f(x) = f(x^*) + \sum_{k=1}^{n-1} \frac{1}{k!} \sum_{i_1, i_2, \dots, i_k = 1}^{m} \frac{\partial^k f(x^*)}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \dots (x_{i_k} - x^*_{i_k}) + \tag{76}$$

$$+ \frac{1}{n!} \sum_{i_1, i_2, \dots, i_n = 1}^{m} \frac{\partial^m f(x_\theta)}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_m}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \dots (x_{i_m} - x^*_{i_m}),$$

where x_θ can be represented, $x_\theta = x^* + \theta(x - x^*)$, $0 < \theta < 1$.

The $k-th$ in the Taylor's theorem can be represented

$$\sum_{i_1, i_2, \dots, i_k = 1}^{m} \frac{f^k(x)}{\partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} (x_{i_1} - x^*_{i_1}) \dots (x_{i_k} - x^*_{i_k}) = D^k f(x)(x - x^*)^{\otimes k}, \tag{77}$$

where $\otimes$ is tensor product, $D : f \to \nabla f$ differentiation operator and $D\otimes D\otimes\ldots\otimes D = D^{\otimes k}$ and $D^{\otimes k} f(x)|_{x=x_\theta} = D^{\otimes k} f(x_\theta)$ is an operater such that $D^{\otimes k} f(x_\theta) : \mathbb{R}^{m^k} \to \mathbb{R}$.
Hence, for $n \geq 4$ expansion (76) can also be represented as

$$\begin{aligned} f(x) =& f(x^*) + Df(x^*)^T (x - x^*) + (x - x^*)^T D^2 f(x^*)(x - x^*) + \\ &+ \sum_{k=3}^{n-1} \frac{1}{k!} D^{\otimes k} f(x^*)(x - x^*)^{\otimes k} + \frac{1}{n!} D^{\otimes n} f(x_\theta)(x - x^*)^{\otimes n}, \end{aligned}$$

where $0 < \theta < 1$.
Since operator $D^{\otimes n} f(x_\theta)$ is finite dimensional, hence it is bounded and we have

$$\left|D^{\otimes n} f(x_\theta)(x - x^*)^{\otimes n}\right| \leq \|D^{\otimes n} f(x_\theta)\| |(x - x^*)^{\otimes n}|. \tag{78}$$

Were we define

$$\|D^{\otimes n} f(x_\theta)\| = \sup_{x\in A} \frac{|D^{\otimes k} f(x_\theta)(x - x^*)^{\otimes k}|}{|(x - x^*)^{\otimes k}|}.$$

which is a norm of the operator. Further, by the definition of the tensor product we have that

$$< x \otimes x, x \otimes x >=< x, x >< x, x >$$

which implies $|x^{\otimes k}| = |x|^k$. Putting that fact together with (78) we get that n-th term in Taylor's Theorem is bounded by

$$\left|D^{\otimes n} f(x_\theta)(x - x^*)^{\otimes n}\right| \leq \|D^{\otimes n} f(x_\theta)\| |x - x^*|^n. \tag{79}$$

If the operator $D^{\otimes k} f(x)$ is invertible, then we have $x = D^{\otimes k} f(x)^{-1} y$ and define

$$\|D^2 f(x)^{-1}\| = \sup_{y\in B} \frac{|D^{\otimes k} f(x)^{-1} y|}{|y|}$$

where B is a range of the considered operator. Then we have inequality

$$\left|D^{\otimes k} f(x)^{-1} y\right| \leq \|D^2 f(x)^{-1}\| |y|,$$

which we transform taking into account relation $x = D^{\otimes k} f(x)^{-1} y$

$$\frac{1}{\|D^2 f(x)^{-1}\|} |x| \leq \left|D^{\otimes k} f(x_\theta) x\right|,$$

setting $x = (x - x^*)^{\otimes k}$ and including $|(x - x^*)^{\otimes k}| = |x - x^*|^k$ we get

$$\frac{|x - x^*|^k}{\|D^2 f(x)^{-1}\|} \leq \left|D^{\otimes k} f(x_\theta)(x - x^*)^{\otimes k}\right|, \tag{80}$$

which is a lower bound for n-th term in the Taylor expansion.
Analogically to Theorem 1 we can define following Taylor's expansion

$$\begin{aligned} \frac{\partial f(x)}{\partial x_j} = \frac{\partial f(x^*)}{\partial x_j} + \sum_{k=1}^{n-1} \frac{1}{k!} \sum_{i_1,i_2,\ldots,i_k=1}^{m} \frac{\partial^{k+1} f(x^*)}{\partial x_j \partial x_{i_1} \partial x_{i_2} \ldots \partial x_{i_k}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \ldots (x_{i_k} - x^*_{i_k}) + \\ + \frac{1}{n!} \sum_{i_1,i_2,\ldots,i_n=1}^{m} \frac{\partial^{m+1} f(x_\theta)}{\partial x_j \partial x_{i_1} \partial x_{i_2} \ldots \partial x_{i_m}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \ldots (x_{i_m} - x^*_{i_m}), \end{aligned} \tag{81}$$

where x_θ is somewhere between x and x^*. The vector composed of elements of above expansion for $j = 1, \dots, m$ can be defined as

$$Df(x) = Df(x^*) + D^2 f(x^*)(x - x^*) + \sum_{k=2}^{n-1} \frac{1}{k!} D^{\otimes(k+1)} f(x^*)(x - x^*)^{\otimes k} + \frac{1}{n!} D^{\otimes(n+1)} f(x_\theta)(x - x^*)^{\otimes n},$$

where we define operator $D^{\otimes(k+1)} f(x) : \mathbb{R}^{m^k} \to \mathbb{R}^m$ for $k \geq 1$ as

$$\left[D^{\otimes(k+1)} f(x)(x - x^*)^{\otimes k}\right]_j = \sum_{i_1, i_2, \dots, i_k = 1}^{m} \frac{\partial^{k+1} f(x)}{\partial x_j \partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \dots (x_{i_k} - x^*_{i_k}),$$

for $j = 1, \dots, m$.
Analogically to the bounds of the remainder of the previous Taylor's expansion (79) and (80) we have for $D^{\otimes(k+1)} f(x)$

$$\left|D^{\otimes(k+1)} f(x_\theta)(x - x^*)^{\otimes k}\right| \leq \|D^{\otimes(k+1)} f(x_\theta)\| |x - x^*|^k. \tag{82}$$

Moreover, if $D^{\otimes(k+1)} f(x)$ is invertible we have

$$\frac{|x - x^*|^k}{\|D^{\otimes(k+1)} f(x_\theta)^{-1}\|} \leq \left|D^{\otimes(k+1)} f(x_\theta)(x - x^*)^{\otimes k}\right|.$$

Further, we define another Taylor's expansion, analogical to one in Theorem 1 as

$$\begin{aligned} \frac{\partial^2 f(x)}{\partial x_j \partial x_l} = \frac{\partial^2 f(x^*)}{\partial x_j \partial x_l} + \sum_{k=1}^{n-1} \frac{1}{k!} \sum_{i_1, i_2, \dots, i_k = 1}^{m} \frac{\partial^{k+2} f(x^*)}{\partial x_j \partial x_l \partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \dots (x_{i_k} - x^*_{i_k}) + \\ + \frac{1}{n!} \sum_{i_1, i_2, \dots, i_n = 1}^{m} \frac{\partial^{m+2} f(x_\theta)}{\partial x_j \partial x_l \partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_m}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \dots (x_{i_m} - x^*_{i_m}), \end{aligned} \tag{83}$$

where again, x_θ is somewhere between x and x^*. The matrix composed of elements of above expansion for $j = 1, \dots, m,\ l = 1, \dots, m$ can be defined as

$$D^2 f(x) = D^2 f(x^*) + \sum_{k=1}^{n-1} \frac{1}{k!} D^{\otimes(k+2)} f(x^*)(x - x^*)^{\otimes k} + \frac{1}{n!} D^{\otimes(n+2)} f(x_\theta)(x - x^*)^{\otimes n},$$

where we define $D^{\otimes(k+2)} f(x) : \mathbb{R}^{m^k} \to \mathbb{R}^{m^2}$ for $k \geq 1$ as

$$\left[D^{\otimes(k+2)} f(x)(x - x^*)^{\otimes k}\right]_{j,l} = \sum_{i_1, i_2, \dots, i_k = 1}^{m} \frac{\partial^{k+2} f(x)}{\partial x_j \partial x_l \partial x_{i_1} \partial x_{i_2} \dots \partial x_{i_k}} (x_{i_1} - x^*_{i_1})(x_{i_2} - x^*_{i_2}) \dots (x_{i_k} - x^*_{i_k}),$$

for $j = 1, \dots, m,\ l = 1, \dots, m$.
Then again we have that

$$\left|D^{\otimes(n+2)} f(x_\theta)(x - x^*)^{\otimes n}\right| \leq \|D^{\otimes(n+2)} f(x_\theta)(x - x^*)^{\otimes n}\| |x - x^*|^n,$$

and if $D^{\otimes(n+2)} f(x_\theta)(x - x^*)^{\otimes n}$ is invertible we have

$$\frac{|x - x^*|^n}{\|D^{\otimes(n+2)} f(x_\theta)(x - x^*)^{\otimes n}\|} \leq \left|D^{\otimes(n+2)} f(x_\theta)(x - x^*)^{\otimes n}\right|,$$

where $n \geq 1$.

Lemma 4. *For any $N > 0$, the integral*

$$\int_{\mathbb{R}^m} e^{-Nx^T Ax} dx = \frac{(2\pi)^{m/2}}{\sqrt{|\det A|}} N^{-\frac{m}{2}} \tag{84}$$

where A is a symmetric negative definite $m-1$-dimensional matrix

Proof. Standard calculus result. ☐

Lemma 5. *For some constant $N > 0$ and $(m-1)$-dimensional sphere with radius R denoted by $S_{m-1}(R)$, with center in the origin, the integral*

$$\int_{\mathbb{R}^m \backslash S_m(R)} e^{Nx^T Ax} dx < \frac{1}{\sqrt{|\det A|}} \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})} e^{-NR^2} \left[\frac{1}{N} \left(\frac{1+\sqrt{1+4R^2}}{2} \right)^{m-1} + \frac{\Gamma(m)}{N^{m-1}} \right], \tag{85}$$

where A is a symmetric negative definite $m-1$-dimensional matrix.

Proof. Since A is negative definite we have

$$\int_{\mathbb{R}^m \backslash S_m(R)} e^{Nx^T Ax} dx = \int_{\mathbb{R}^m \backslash S_m(R)} e^{-N|x^T Ax|} dx$$

We start we performing the change of variables $x = \sqrt{A}x'$ and we get

$$\int_{\mathbb{R}^m \backslash S_{m-1}(R)} e^{-N|x^T Ax|} dx = \frac{1}{\sqrt{|\det A|}} \int_{\mathbb{R}^m \backslash S_m(R)} e^{-N|x|^2} dx. \tag{86}$$

Then we change the coordinate system into spherical one where $x' = rs_r$, and radius $r = |x'|$

$$\int_{\mathbb{R}^m \backslash S_m(R)} e^{-N|x|^2} dy = \int_R^\infty \int_{S_m(r)} e^{-Nr^2} ds_r dr$$

As the function under the integral does not depend on the surface coordinates we can independently integrate over the surface. The surface of sphere in $m-1$-dimensional space of radius r is given by

$$S_m(r) = 2\frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})} r^{m-2},$$

and then

$$\int_R^\infty \int_{S_m(r)} e^{-Nr^2} ds_r dr = 2\frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})} \int_R^\infty r^{m-1} e^{-Nr^2} dr.$$

Now we perform a substitution $t = N(r^2 - R^2)$ and get

$$\int_R^\infty r^{m-1} e^{-Nr^2} dr = e^{-NR^2} \int_0^\infty e^{-t} \left(\frac{t}{N} + R^2 \right)^{\frac{m-1}{2}} \frac{dt}{2N}. \tag{87}$$

For $t > 0$ we have following estimate

$$\left(\frac{t}{N} + R^2 \right)^{\frac{m-1}{2}} < \left(\frac{1+\sqrt{1+4R^2}}{2} \right)^{m-1} + \left(\frac{t}{N} \right)^{m-1}$$

where K is some positive constant. We substitute above into (87) and get

$$\int_R^\infty r^{m-1}e^{-Nr^2}dr < e^{-NR^2}\int_0^\infty e^{-t}\left(\frac{1+\sqrt{1+4R^2}}{2}\right)^{m-1}\frac{dt}{2N} + \int_0^\infty e^{-t}K\left(\frac{t}{N}\right)^{m-1}\frac{dt}{2N}.$$

Then after calculating explicitely the first integral and noting that gamma integral representation of gamma function is

$$\Gamma(z) = \int_0^\infty t^{z-1}e^{-t}dt,$$

we obtain

$$\int_R^\infty r^{m-1}e^{-Nr^2}dr < e^{-NR^2}\frac{1}{2N}\left(\frac{1+\sqrt{1+4R^2}}{2}\right)^{m-1} + e^{-NR^2}\frac{K}{2N^m}\Gamma(m).$$

Now we put above together with (A), (86), (A)

$$\int_{\mathbb{R}^m\setminus S_m(R)} e^{Nx^TAx}dx < \frac{1}{\sqrt{|\det A|}}\frac{\pi^{\frac{m-1}{2}}}{\Gamma(\frac{m}{2})}e^{-NR^2}\left[\frac{1}{N}\left(\frac{1+\sqrt{1+4R^2}}{2}\right)^{m-1} + \frac{\Gamma(m-1)}{N^m}\right],$$

hence we get the final result. ☐

Lemma 6. *For some positive a and an integer m we have following estimate valid for all $t \geq 0$*

$$(t+a)^{m/2} < \left(\frac{1-\sqrt{1+4a}}{2}\right)^m + t^m \tag{88}$$

Proof. Let us calculate the points of intersection of functions $(t+a)^{m/2}$ and t^m. For this we have to solve

$$(t+a)^{m/2} = t^m,$$

and this is equivalent to quadratic equation

$$t^2 - t - a = 0,$$

which has two roots

$$t_1 = \frac{1+\sqrt{1+4a}}{2},\ t_2 = \frac{1-\sqrt{1+4a}}{2},$$

and since $a > 0$ second root $t_2 < 0$. We consider $t \geq 0$ and as both functions are strictly increasing for $t < t_1$ function $(t+a)^{m/2}$ is bigger than t^m and for $y > y_1$ other way round. Hence by adding to the function t^m a constant $(t_1)^m$ we obtain situation when it is bigger than $(t+a)^{m/2}$ for all $t \geq 0$. ☐

Theorem 5 (Matrix series). *For the matrix $A \in \mathbb{R}^{m\times m}$ we have following exapnsions*

$$e^A = 1 + A + \frac{A^2}{2!} + \frac{A^3}{3!} + \dots, \tag{89}$$

$$(I+A)^{-1} = 1 - A + A^2 - A^3 + \dots, \tag{90}$$

$$\log(I+A) = A - \frac{A^2}{2} + \frac{A^3}{3} - \dots, \tag{91}$$

where for the last two series we have conditions that $||A|| < 1$.

Theorem 6 (Determinant approximation)**.** *For a matrix* $A \in \mathbb{R}^{m\times m}$ *we have following approximation*

$$\det\left(I+\frac{A}{N}\right) = 1+\frac{tr(A)}{N}+O\biggl(\frac{1}{N^2}\biggr), \tag{92}$$

as $N \to \infty$*.*

Proof. From the Jacobi formula we have that

$$\det\left(I+\frac{A}{N}\right) = \exp(\mathrm{tr}(\log\left(I+\frac{A}{N}\right))). \tag{93}$$

for any A.
From the expansion of the logarithm and exponential of a matrix form previous thereom we have that

$$e^{\frac{A}{N}} = 1+\frac{A}{N}+O\biggl(\frac{1}{N^2}\biggr),$$
$$\log\left(I+\frac{A}{N}\right) = \frac{A}{N}+O\biggl(\frac{1}{N^2}\biggr),$$

when $N \to \infty$. Substituting above asymptotic equations into (93) we get

$$\det\left(I+\frac{A}{N}\right) = 1+\mathrm{tr}\biggl(\frac{A}{N}+O\biggl(\frac{1}{N^2}\biggr)\biggr)+O\biggl(\frac{1}{N^2}\biggr), \tag{94}$$

for $N \to \infty$.
Since trace is a linear transformation above is equal to the equation from the theorem. ☐

B Asymptotic theory and approximation

Let $f : A \to \mathbb{R}$ be a continuous function and $A = (a, \infty)$ for some a.

Definition 1 (Big O)**.** *The function* f *is of order* O *of the function* $g : A \to \mathbb{R}$ *as* $x \to \infty$ *if there exists is a constant* $K > 0$ *and* $x_K \in A$ *such that for all* $x > x_K$

$$|f(x)| \leq K|g(x)|,$$

and we write it symbolically

$$f(x) = O(g(x)),\ x \to \infty.$$

Definition 2 (Small o)**.** *The function* f *is of order* o *of the function* $g : A \to \mathbb{R}$ *as* $x \to \infty$ *if for all* $K > 0$ *there exists* $x_K \in A$ *such that for all* $x > x_K$

$$|f(x)| \leq K|g(x)|,$$

and we write it symbolically

$$f(x) = o(g(x)),\ x \to \infty.$$

Lemma 7. *Given inequalities*

$$A_i - B_i = C_i, \;\; i = 1, \ldots, m, \tag{95}$$
$$\tag{96}$$

where $m \in \mathbb{N}$ *following inequalities holds*

$$\prod_{i=1}^{m} A_i - \prod_{i=1}^{m} B_i = \prod_{i=1}^{m} C_i + \sum_{j=1}^{m-1} \sum_{C_m^{m-j,j}} \prod_{k=1}^{j} \prod_{l=j+1}^{m} C_{i_k} B_{i_l},$$

Proof. We start by introducing equality

$$\prod_{i=1}^{m} A_i = (A_1 - B_1) \prod_{i=2}^{m} A_i + B_1 \prod_{i=2}^{m} A_i,$$

which we obtained by adding and deducting $B_1 \prod_{i=2}^{m} A_i$ to $\prod_{i=1}^{m} A_i$.
Then again, we add and deduce, but this time $B_1 B_2 \prod_{l=3}^{m} A_l$ and $(A_1 - B_1) B_2 \prod_{l=3}^{m} A_l$ and get

$$\prod_{i=1}^{m} A_i = (A_1 - B_1)(A_2 - B_2) \prod_{i=3}^{m} A_i + (A_1 - B_1) B_2 \prod_{i=3}^{m} A_i + B_1 (A_2 - B_2) \prod_{i=3}^{m} A_i + B_1 B_2 \prod_{i=3}^{m} A_i.$$

We repeat that step until all A_i's in the products are replaced by $(A_i - B_i)$, which eventually leads to the equation

$$\prod_{i=1}^{m} A_i - \prod_{i=1}^{m} B_i = \prod_{i=1}^{m} (A_i - B_i) + \sum_{j=1}^{m-1} \sum_{C_m^{m-j,j}} \prod_{k=1}^{j} \prod_{l=j+1}^{m} (A_{i_k} - B_{i_k}) B_{i_l} \tag{97}$$

where $\sum_{C_m^{m-j,j}}$ is a sum over possible arrangements of the elements of the set $\{1, 2, \ldots, m\}$ into two groups, where elements does not repeat and within the group the order does not matter. First group is of the size $m - j$ and second j and their elements correspond, respectively, to the indecies $i_k, k = 1, \ldots, m - j$ and $i_l, l = m - j + 1 \ldots, m$.
Then we substitute (95) and obtain

$$\prod_{i=1}^{m} A_i - \prod_{i=1}^{m} B_i = \prod_{i=1}^{m} C_i + \sum_{j=1}^{m-1} \sum_{C_m^{m-j,j}} \prod_{k=1}^{j} \prod_{l=j+1}^{m} C_{i_k} B_{i_l},$$

which is our result. □

Lemma 8. *Given equalities*

$$A_1 - B_1 = \sigma_1, \tag{98}$$
$$A_2 - B_2 = \sigma_2, \tag{99}$$

where $\sigma_1, \sigma_2 \in \mathbb{R}$ *are some constants, following equality holds*

$$\frac{A_1}{A_2} - \frac{B_1}{B_2} = -\frac{B_1 \sigma_2}{B_2 (B_2 \sigma_2)} + \frac{\sigma_1}{B_2 + \sigma_2}.$$

Proof. First we divide (98) by (99) and get

$$\frac{A_1}{A_2} = \frac{B_1 + \sigma_1}{B_2 - \sigma_2}, \tag{100}$$

and then we transform LHS

$$\frac{B_1 + \sigma_1}{B_2 + \sigma_2} = \frac{B_1}{B_2 + \sigma_2} + \frac{\sigma_1}{B_2 + \sigma_2},$$

then add and deduce B_1/B_2 and perform some manipulations

$$\begin{aligned} \frac{B_1}{B_2 + \sigma_2} + \frac{\sigma_1}{B_2 + \sigma_2} &= \frac{B_1}{B_2} + \frac{B_1}{B_2 + \sigma_2} - \frac{B_1}{B_2} + \frac{\sigma_1}{B_2 + \sigma_2} = \\ &= \frac{B_1}{B_2} + \frac{B_1 B_2}{B_2(B_2 + \sigma_2)} - \frac{B_1(B_2 + \sigma_2)}{B_2(B_2 + \sigma_2)} + \frac{\sigma_1}{B_2 + \sigma_2} = \\ &= \frac{B_1}{B_2} - \frac{B_1 \sigma_2}{B_2(B_2 + \sigma_2)} + \frac{\sigma_1}{B_2 + \sigma_2}, \end{aligned}$$

and then we put it back to (100), change the side of B_1/B_2 and obtain final result

$$\frac{A_1}{A_2} - \frac{B_1}{B_2} = -\frac{B_1 \sigma_2}{B_2(B_2 + \sigma_2)} + \frac{\sigma_1}{B_2 + \sigma_2}.$$

□

References

Vassili N. Kolokoltsov. *Semiclassical Analysis for Diffusions and Strochastic processes.* Springer, 2000.